

# 1:2 Lead(2+) salts of long-chain carboxylic acids: Environment tier II assessment

04 July 2014

**CAS Registry Numbers: 1072-35-1, 58405-97-3, 15773-56-5, 94266-31-6, 29597-84-0, 93858-24-3, 13094-04-7, 1120-46-3, 33627-12-2.**



- Preface
- Disclaimer
- Grouping Rationale
- Chemical Identity
- Physical and Chemical Properties
- Import, Manufacture and Use
- Environmental Regulatory Status
- Environmental Exposure
- Environmental Effects
- Categorisation of Environmental Hazard
- Risk Characterisation
- Key Findings
- Recommendations
- Environmental Hazard Classification
- References

## Preface

This assessment was carried out by staff of the National Industrial Chemicals Notification and Assessment Scheme (NICNAS) using the Inventory Multi-tiered Assessment and Prioritisation (IMAP) framework.

The IMAP framework addresses the human health and environmental impacts of previously unassessed industrial chemicals listed on the Australian Inventory of Chemical Substances (the Inventory).

The framework was developed with significant input from stakeholders and provides a more rapid, flexible and transparent approach for the assessment of chemicals listed on the Inventory.

Stage One of the implementation of this framework, which lasted four years from 1 July 2012, examined 3000 chemicals meeting characteristics identified by stakeholders as needing priority assessment. This included chemicals for which NICNAS already held exposure information, chemicals identified as a concern or for which regulatory action had been taken overseas, and chemicals detected in international studies analysing chemicals present in babies' umbilical cord blood.

Stage Two of IMAP began in July 2016. We are continuing to assess chemicals on the Inventory, including chemicals identified as a concern for which action has been taken overseas and chemicals that can be rapidly identified and assessed by using Stage One information. We are also continuing to publish information for chemicals on the Inventory that pose a low risk to human health or the environment or both. This work provides efficiencies and enables us to identify higher risk chemicals requiring assessment.

The IMAP framework is a science and risk-based model designed to align the assessment effort with the human health and environmental impacts of chemicals. It has three tiers of assessment, with the assessment effort increasing with each tier. The Tier I assessment is a high throughput approach using tabulated electronic data. The Tier II assessment is an evaluation of risk on a substance-by-substance or chemical category-by-category basis. Tier III assessments are conducted to address specific concerns that could not be resolved during the Tier II assessment.

These assessments are carried out by staff employed by the Australian Government Department of Health and the Australian Government Department of the Environment and Energy. The human health and environment risk assessments are conducted and published separately, using information available at the time, and may be undertaken at different tiers.

This chemical or group of chemicals are being assessed at Tier II because the Tier I assessment indicated that it needed further investigation.

For more detail on this program please visit: [www.nicnas.gov.au](http://www.nicnas.gov.au).

## Disclaimer

NICNAS has made every effort to assure the quality of information available in this report. However, before relying on it for a specific purpose, users should obtain advice relevant to their particular circumstances. This report has been prepared by NICNAS using a range of sources, including information from databases maintained by third parties, which include data supplied by industry. NICNAS has not verified and cannot guarantee the correctness of all information obtained from those databases. Reproduction or further distribution of this information may be subject to copyright protection. Use of this information without obtaining the permission from the owner(s) of the respective information might violate the rights of the owner. NICNAS does not take any responsibility whatsoever for any copyright or other infringements that may be caused by using this information.

### Acronyms & Abbreviations

## Grouping Rationale

This Tier II assessment considers the environmental risks associated with industrial uses of the lead(2+) salts of nine closely related carboxylic acids:


- Octadecanoic acid, lead(2+) salt (Lead distearate)
- Octadecanoic acid, 12-hydroxy-, lead(2+) salt (2:1) (Lead bis(12-hydroxystearate))
- Hexadecanoic acid, lead(2+) salt (Lead dipalmitate)
- Eicosanoic acid, lead(2+) salt (2:1) (Lead diicosanoate)
- Docosanoic acid, lead(2+) salt (Lead dibehenate)
- 9-Hexadecenoic acid, lead(2+) salt, (Z)- (Lead dipalmitoleate)
- 9-Octadecenoic acid, 12-hydroxy-, lead(2+) salt (2:1), [R-(Z)] (Lead diricinoleate)
- 9-Octadecenoic acid, lead(2+) salt, (Z)- (Lead dioleate)
- 9,12-Octadecadienoic acid, (Z,Z)-, lead(2+) salt (Lead dilinoleate)


The parent carboxylic acids for these salts are all long-chain (greater than C<sub>12</sub>) naturally occurring saturated, mono-unsaturated, di-unsaturated or hydroxylated fatty acids. The lead(2+) salts of these acids are a subset of a large class of industrially important metal salts of fatty acids collectively identified as metallic soaps. This subset of metallic soaps is of definite chemical composition based on their Chemical Abstracts Service chemical names and formulae as listed in the Australian Inventory of Chemical Substances (the Inventory). The entries on the Inventory specify the ratio of lead(2+) to carboxylic acid in these salts as 1:2.


The risk assessment of these chemicals has been conducted as a group because all nine substances are well-defined lead salts of structurally and chemically similar long-chain fatty acids. Although these salts are sparingly soluble in water, they can all potentially release toxic lead(2+) ions which provides a common source of toxicity for each of these substances. Furthermore, the potential environmental exposure pathways for chemicals in this group are similar based on specific industrial uses as stabilisers in polyvinyl chloride (PVC) plastics, as well as potential uses in paints, coatings and greases. Based on these considerations, the chemicals in this group are each expected to have generally similar environmental fate and ecotoxicity profiles.

The chemicals in this group are related to the chemicals assessed as part of the Lead Salts of Long-Chain Carboxylic Acids group. Both groups of chemicals are structurally similar, but they may have some minor compositional differences. Although they have been assessed as separate groups, persons interested in the chemicals assessed here should also consider those assessed under the Lead Salts of Long-Chain Carboxylic Acids group assessment.


## Chemical Identity

CAS RN	1072-35-1
Chemical Name	Octadecanoic acid, lead(2+) salt
Synonyms	Lead distearate Lead distearate, pure Neutral lead distearate Lead(2+) dioctadecanoate
Structural Formula	
Molecular Formula	C <sub>36</sub> H <sub>70</sub> O <sub>4</sub> Pb
Molecular Weight (g/mol)	774.14
SMILES	[Pb++].CCCCCCCCCCCCCCCC(=O) [O-].CCCCCCCCCCCCCCCC(=O)[O-]


CAS RN	58405-97-3
Chemical Name	Octadecanoic acid, 12-hydroxy-, lead(2+) salt (2:1)
Synonyms	Lead bis(12-hydroxystearate) Lead(2+) bis(12-hydroxyoctadecanoate)
Structural Formula	
Molecular Formula	C <sub>36</sub> H <sub>70</sub> O <sub>6</sub> Pb
Molecular Weight (g/mol)	806.14
SMILES	[Pb++].CCCCCCC(O)CCCCCCCCCCCC(=O) [O-].CCCCCCC(O)CCCCCCCCCCCC(=O)[O-]

CAS RN	15773-56-5
Chemical Name	Hexadecanoic acid, lead(2+) salt
Synonyms	Lead dipalmitate Lead(2+) dihexadecanoate
Structural Formula	


Molecular Formula	$C_{32}H_{62}O_4Pb$
Molecular Weight (g/mol)	718.03
SMILES	[Pb++].CCCCCCCCCCCCCCCC(=O)[O-].CCCCCCCCCCCCCCCC(=O)[O-]

CAS RN	94266-31-6
Chemical Name	Eicosanoic acid, lead(2+) salt (2:1)
Synonyms	Lead diicosanoate Lead icosanoate (1:2) Lead(2+) diicosanoate
Structural Formula	
Molecular Formula	$C_{40}H_{78}O_4Pb$
Molecular Weight (g/mol)	830.25
SMILES	[Pb++].CCCCCCCCCCCCCCCCCCCC(=O)[O-].CCCCCCCCCCCCCCCCCCCC(=O)[O-]


CAS RN	29597-84-0
--------	------------

Chemical Name	Docosanoic acid, lead(2+) salt
Synonyms	Lead dibehenate Lead(2+) didocosanoate
Structural Formula	
Molecular Formula	C <sub>44</sub> H <sub>86</sub> O <sub>4</sub> Pb
Molecular Weight (g/mol)	886.35
SMILES	[Pb <sup>++</sup> ].CCCCCCCCCCCCCCCCCCCCCCCC(=O) [O-].CCCCCCCCCCCCCCCCCCCCCCCC(=O)[O-]


CAS RN	93858-24-3
Chemical Name	9-Hexadecenoic acid, lead(2+) salt, (Z)-
Synonyms	Lead dipalmitoleate Lead(2+) di[(9Z)-hexadec-9-enoate]
Structural Formula	



Molecular Formula	C <sub>32</sub> H <sub>58</sub> O <sub>4</sub> Pb
Molecular Weight (g/mol)	714.00
SMILES	[Pb++].CCCCC/C=C\CCCCCCC(=O) [O-].CCCCC/C=C\CCCCCCC(=O)[O-]
CAS RN	13094-04-7
Chemical Name	9-Octadecenoic acid, 12-hydroxy-, lead(2+) salt (2:1), [ <i>R</i> -( <i>Z</i> )]
Synonyms	Lead diricinoleate Lead(2+) bis[(9 <i>Z</i> ,12 <i>R</i> )-12-hydroxyoctadec-9-enoate]
Structural Formula	



Molecular Formula	C <sub>36</sub> H <sub>66</sub> O <sub>6</sub> Pb
Molecular Weight (g/mol)	802.11
SMILES	[Pb++].CCCCC[C@@]([H])(O)C/C=C\CCCCCCCC(=O)[O-].CCCCC[C@@]([H])(O)C/C=C\CCCCCCCC(=O)[O-]
CAS RN	1120-46-3
Chemical Name	9-Octadecenoic acid, lead(2+) salt, (Z)-
Synonyms	Lead dioleate Lead(2+) di[(9Z)-octadec-9-enoate]
Structural Formula	



Molecular Formula	C <sub>36</sub> H <sub>66</sub> O <sub>4</sub> Pb
Molecular Weight (g/mol)	770.11
SMILES	[Pb++].CCCCCCCC/C=C\CCCCCCCC(=O) [O-].CCCCCCCC/C=C\CCCCCCCC(=O)[O-]
CAS RN	33627-12-2
Chemical Name	9,12-Octadecadienoic acid, (Z,Z)-, lead(2+) salt
Synonyms	Lead dilinoleate Lead(2+) di[(9Z,12Z)-octadeca-9,12-dienoate]
Structural Formula	

Molecular Formula	C <sub>36</sub> H <sub>62</sub> O <sub>4</sub> Pb
Molecular Weight (g/mol)	766.08
SMILES	[Pb++].CCCCC/C=C\C/C=C\C\CCCCCCCC(=O) [O-].CCCCC/C=C\C/C=C\C\CCCCCCCC(=O)[O-]

## Physical and Chemical Properties

Lead distearate is a white solid with a melting point of 125°C (LMC, 2013; Nora and Koenen, 2012). The commercially produced substance has a free fatty acid content of 1% w/w and a total lead content of 27-28% w/w. Limited physical property data are available for the remaining chemicals in this group. However, the lead(2+) salts of other fatty acids in this group are all expected to be solids under ambient conditions based on the typical properties of metallic soaps (Nora and Koenen, 2012).

The metallic soaps of divalent (2+) metal ions are well known to be only sparingly soluble in water (Nora and Koenen, 2012). A number of studies of the solubility of various divalent metallic soaps show the lead(2+) soaps typically dissociate in water to release lead(2+) ions and fatty acids in the form of carboxylate mono-anions. These studies also clearly demonstrate that the water solubility of metallic soaps decreases as the length of the fatty acid carbon chain increases. Additionally, these studies have shown that the lead(2+) salts are typically the least soluble of the divalent metal salts of fatty acids (Hunter and Liss, 1976; Mauchauffee, et al., 2008 ).

Solubility products ( $K_{sp}$ s), which are a quantitative measure of the position of the solubility equilibrium, are available for lead dipalmitate and lead distearate (Hunter and Liss, 1976). The measurements, obtained at 25°C and an ionic strength of 0.1 M (NaCl), are presented in the following table. The concentrations of lead(2+) ions at saturation were calculated using an activity coefficient of 0.586 M, as used in the original study:

Chemical	Lead dipalmitate	Lead distearate
$K_{sp}$	$6.31 \times 10^{-22}$	$1.99 \times 10^{-24}$
Lead(2+) Concentration at Saturation	$20 \times 10^{-3}$ mg/L	$2.84 \times 10^{-3}$ mg/L

No quantitative water solubility data have been identified for lead diicosanoate (di-C<sub>20</sub>) and lead dibehenate (di-C<sub>22</sub>). However, both of these salts are expected to be less soluble than lead distearate (di-C<sub>18</sub>) based on the observed trend in solubility for metallic soaps.

Further, no quantitative water solubility data have been identified for the other salts in this group. Based on the measured solubility products for the salts of saturated fatty acid analogues, it can be assumed that the salts of the other fatty acids in this group are also sparingly soluble in water. However, the lead(2+) concentration at saturation cannot be estimated for these salts.

## Import, Manufacture and Use

### Australia

Lead distearate has been reported to be used in Australia predominantly in the production of non-pressurised PVC pipe products as an additive and stabiliser. The chemical was listed on the 2002 High Volume Industrial Chemical List (HVICL) with a reported annual introduction volume between 1000 and 9999 tonnes. However, the chemical is not listed on the most recent HVICL, published in 2006 (EPT, 2003; NICNAS, 2006; 2013; 2014).

In 2012, the Vinyl Council Australia announced that all signatories to their Product Stewardship Program had completed a phase out of lead-based stabiliser compounds in PVC products (Vinyl Council Australia, 2012). The sole manufacturer of PVC resin in Australia, which supplies two thirds of domestic demand, is a signatory to the program. In addition, several PVC product manufacturers, key additives suppliers and PVC importers are also signatories (Chemiplas, 2012; Tilley, 2013; Vinyl Council Australia, 2014).

No specific Australian use, import or manufacturing information has been identified for the remaining chemicals in this group.

### International

Available information shows that lead(2+) salts of long-chain fatty acids were widely used historically as additives in extreme pressure greases (Calhoun, 1960; OECD, 2004; Shugarman, 2003), with some information also suggesting use as driers in paint (Schwartz, 2012). Further available information indicates that lead distearate has recent reported industrial use as a stabiliser in the manufacture of PVC products (Health Canada, 2013; Nordic Council of Ministers, 2013; PVC Europe, 2014 ).

Lead distearate was reported to be used in some Nordic countries between 2009 and 2011 (Nordic Council of Ministers, 2013). However, it is noted that lead distearate – alongside all chemicals in this group – is only pre-registered under the Registration, Evaluation, Authorisation and Restriction of Chemicals (REACH) legislation (ECHA, 2014b). Therefore, it can be assumed that these chemicals are not currently being used at a volume greater than 100 tonnes per annum in the European Union (ECHA, 2014c).

The European Stabiliser Producers Association has reported that all lead-based PVC stabilisers with current commercial relevance have undergone the complete registration process under the REACH legislation (ESPA, 2014). Therefore, it can be assumed that the chemicals in this group are not currently considered to be commercially relevant PVC stabilisers in the European Union. Examples of lead-based PVC stabilisers that are registered include a mixture of C<sub>16-18</sub> lead carboxylates (CAS RN 91031-62-8) and dibasic lead stearate (CAS RN 12578-12-0) (ECHA, 2014d; 2014e). Nevertheless, use of lead-based PVC stabilisers is declining rapidly in the European Union, with the European PVC industry agreeing to completely phase out use by 2015. As a result of the agreement, lead-based PVC stabiliser use decreased by more than 75% between 2000 and 2010, exceeding the milestone targets that had been set (Vinyl 2010, 2006; 2011 ).

No additional use data were available for the chemicals in this group. Apart from lead distearate, the chemicals in this group do not appear to be used industrially in Canada and have not been introduced for an industrial use at a volume greater than 100 kg per annum in Sweden since its records began in 1992 (Environment Canada, 2013b; Keml, 2014).

## Environmental Regulatory Status

### Australia

Lead and lead compounds are subject to reporting under the Australian National Pollutant Inventory (NPI). Under the NPI, emissions of lead and lead compounds are required to be reported annually by facilities that use or emit more than 10 tonnes of lead or lead compounds, burn more than 2000 tonnes of fuel, consume more than 60 000 megawatt hours of electricity (excluding lighting and motive purposes), or have an electricity rating of 20 megawatts during a reporting year (Australian

Government Department of the Environment, 2013b). Emissions may be intentional, accidental or incidental releases arising through industrial processes. Additionally, emissions of lead and lead compounds from diffuse sources, such as lawn mowers and wood heaters, are also periodically estimated by state environment authorities. Diffuse emissions data are updated much less frequently than facility data (Australian Government Department of the Environment, 2013a).

The *Uniform Paint Standard* of the *Poisons Standard 2013*(Cwlth) recommends that lead and lead compounds be limited to 0.1% of the non-volatile content of all paints manufactured and sold in Australia (measured as elemental lead) (Commonwealth of Australia, 2013).

## United Nations

No chemicals in this group are currently identified as a Persistent Organic Pollutant (UNEP, 2001), ozone depleting substance (UNEP, 1985), or hazardous substance for the purpose of international trade (UNEP & FAO, 1998).

The United Nations Environment Programme (UNEP), in partnership with the World Health Organization (WHO), has established the Global Alliance to Eliminate Lead Paint. The alliance primarily supports actions in developing nations to reduce and phase-out lead use in paints (UNEP, 2011).

## OECD

No chemicals in this group have been sponsored for assessment under the Cooperative Chemicals Assessment Programme (CoCAP). However, lead distearate has been identified as a High Production Volume chemical by the OECD. High Production Volume chemicals are used at more than 1000 tonnes per annum in at least one member country (OECD, 2012; 2013).

## Canada

Lead distearate was categorised as Persistent (P), not Bioaccumulative (not B), and Inherently Toxic to the Environment (iT<sub>E</sub>) by Environment Canada during the Categorization of the Domestic Substances List (DSL) (Environment Canada, 2013c). A subsequent screening assessment is yet to be completed (Environment Canada, 2013a).

The remaining chemicals in this group are not currently listed on the DSL (Environment Canada, 2013b).

Lead in most paint products available to consumers in Canada is limited to 90 mg/kg (0.009% w/w) of the non-volatile content under the *Surface Coating Materials Regulations* of the *Canada Consumer Product Safety Act 2010*. Exempted products which contain more than 90 mg/kg lead must be appropriately labelled (Government of Canada, 2005).

## European Union

All chemicals in this group have been pre-registered, but have not yet undergone the full registration process, under the REACH legislation (ECHA, 2014a). Therefore, no chemicals in this group are currently identified as a Substance of Very High Concern in the European Union.

However, it is noted that all other lead compounds currently registered under the REACH legislation (for example, mixed C<sub>16-18</sub> lead carboxylates (CAS RN 91031-62-8)) have been identified as Substances of Very High Concern due to their reproductive toxicity. These chemicals are all currently included on the Candidate List for Eventual Inclusion in Annex XIV (the Authorisation List) to REACH (ECHA, 2013).

## United States of America

Select industries in the United States of America (USA), including most mining, utilities and manufacturing facilities, must report emissions of lead and lead compounds to the United States Environmental Protection Agency (US EPA) if the facility manufactures or processes more than 25 000 pounds of lead compounds (approximately 11 tonnes) per annum. Reporting is

also required if the facility otherwise uses more than 10 000 pounds (approximately 4.5 tonnes) in a reporting year (US EPA, 2012; 2014a; 2014b).

Most paint products containing lead, as well as toys and furniture painted using such products, are prohibited in the USA if the lead content exceeds 0.009% w/w of the non-volatile paint content under the *Consumer Product Safety Improvement Act 2008*. Exempted products which contain more than 0.009% w/w lead must be appropriately labelled (United States Government, 2008).

## Environmental Exposure

The expected principal industrial use of lead distearate is as a stabiliser in the manufacture of PVC articles, at a typical concentration of approximately 0.5% (Smith, 1998). This use pattern has limited potential for direct release of lead distearate to the environment because the substance is contained within the insoluble polymer matrix of durable articles. However, over time, and depending on the use of the PVC article, there is the potential for release of some of the lead(2+) ions from the stabiliser in the form of soluble ionic lead compounds (Al-Malack, 2001; Koh, et al., 1991; Lasheen, et al., 2008). These soluble lead compounds can be released to the terrestrial compartment or aquatic compartment depending on the situation under which the PVC article is used. The major proportion of the lead distearate that is incorporated into a PVC matrix is expected to be retained within the manufactured article and ultimately disposed of to landfill.

The extent to which the remaining chemicals in this group are used in Australia is unclear. Based on the use information available for lead distearate, international data and the chemical similarity of the substances addressed in this group assessment, the remaining chemicals may have some speciality uses, potentially including use as PVC additives, as driers in paints and coatings, and in greases. However, the chemicals in this group were not identified as being used in paints in a survey conducted by the Australian Paint Manufacturers' Federation Inc. and NICNAS in 2007 (NICNAS, 2007), and available international data suggests current use in greases is unlikely (Shugarman, 2003).

For the purposes of this assessment, it will be assumed that any current domestic industrial uses of the other lead(2+) salts of long-chain fatty acids in this group will be similar to those identified for lead distearate. Direct releases of these salts to the environment are therefore considered unlikely. However, diffuse emissions of lead(2+) ions from PVC articles may occur.

## Environmental Fate

### Dissolution, Speciation and Partitioning

The behaviour of the lead(2+) ion is strongly dependent on the chemistry of the environmental compartment into which it is released.

The metallic soaps are salts of weak acids and dissolution of these substances in water therefore results in hydrolysis. Hydrolysis of the fatty acid conjugate base results in the formation of the corresponding fatty acid. The extent to which this reaction occurs is dependent on the properties of the acid and the pH of the aquatic environment. At neutral pH, fatty acids are predominantly in the conjugate base (anionic) form. For the metal ion, hydrolysis is a result of dissociation of hydrogen ions from water molecules coordinated to the metal ion. In the case of lead(2+) ions, coordinated water is only weakly acidic and the predominant form of the dissolved metal ion in pure water at neutral pH is the simple hydrated lead(2+) di-cation (Mauchauffee, et al., 2008).

However, in environmental waters, lead(2+) ions are expected to undergo a range of additional reactions and they can therefore exist as a variety of different chemical species. The speciation chemistry of lead(2+) in environmental waters is complex and strongly dependent on water chemistry, especially pH and the presence of complexing anions. For example, at pH levels lower than 7, the simple hydrated lead(2+) di-cation is the dominant species of ionic lead (ANZECC, 2000a). However, at pH levels greater than 7, the speciation of ionic lead is dominated by the formation of carbonate complexes, such as  $PbCO_3$  (US EPA, 1999). These carbonate complexes have low solubility and the precipitation of lead(2+) salts becomes important under alkaline conditions. In seawater, lead(2+) is complexed by chloride ions. These complexes account for more than 90% of the speciation of ionic lead in seawater (ANZECC, 2000a).

In addition to complexation by simple inorganic anions, lead(2+) is complexed by organic matter in environmental waters. These complexes are of great significance for the environmental fate of lead(2+) because they account for the majority of dissolved lead in natural freshwater at pH values in the range 5 to 9 (ANZECC, 2000a). Further, the partitioning of lead(2+) from the water phase onto clay particles, particulate humic substances, and iron, aluminium and manganese (oxy)hydroxides have all been identified as important mechanisms controlling the concentration of lead in natural waters. In sediments, the speciation of lead will be dominated by complexes with humic substances and by the formation of relatively insoluble lead sulfides (ANZECC, 2000a).

The partitioning behaviour of lead in soil has a complex dependence on soil properties, but is mainly determined by pH and clay content. A detailed review of measured partition coefficients for sorption from water onto soil ( $K_{Ds}$ ) conducted by the US EPA identified values for lead(2+) in the range of 150 L/kg to 44 580 L/kg. The adsorption of lead on soil generally increases with increasing soil pH (US EPA, 1999). Lead is typically considered relatively immobile in soil (UNEP, 2010).

## Biotransformation

The organic components of the salts in this group are biodegradable both as the lead(2+) salts and as the free fatty acids.

A ready biodegradability study on lead distearate, conducted in accordance with OECD Test Guideline (TG) 301C, found the chemical to undergo 32.9% degradation in 14 days. The predominant degradation products were lead dipalmitate (di-C<sub>16</sub>) and lead myristate (C<sub>14</sub>) (LMC, 2013; NITE, 2014). These findings suggest that the aliphatic chains of the chemicals in this group are available to undergo a degree of biodegradation despite the relatively low solubility of the parent lead(2+) salt in water. It should be noted that as the aliphatic chain of these chemicals is shortened, the product lead(2+) salts will become relatively more soluble, which could increase the bioavailability of the ionic lead component in the environment.

The fatty acid components of the chemicals in this group are ubiquitous, naturally occurring dietary fats that will undergo relatively rapid biodegradation in the environment. Available experimental data from studies conducted in accordance with standard test guidelines demonstrate that linoleic acid, oleic acid, palmitic acid and 12-hydroxystearic acid are all readily biodegradable (80% degradation in 28 days, OECD TG 301C; 78% degradation in 28 days, OECD TG 301C, 65% degradation in 28 days, ISO 10708 and 93% degradation in 28 days, OECD TG 301C, respectively) (LMC, 2013). The two unsaturated fatty acids, palmitoleic acid and ricinoleic acid, are also expected to be rapidly biodegradable in the environment based on their structural similarity to the other unsaturated fatty acids for which biodegradation data are available.

The saturated fatty acids with longer aliphatic chains experience less rapid biodegradation in laboratory tests. For example, although 71% of stearic acid degraded in 28 days in a study conducted in accordance with OECD TG 301B, the "10 day window" requirement was not met and the chemical is therefore not classed as readily biodegradable (LMC, 2013). Behenic acid is not readily biodegradable based on a study conducted in accordance with OECD TG 301C (52% degradation in 28 days), but is inherently biodegradable according to a study conducted in accordance with OECD TG 302C (74% degradation in 28 days) (LMC, 2013). Eicosanoic acid (C<sub>20</sub>), which has a structure intermediate between that of stearic acid (C<sub>18</sub>) and behenic acid (C<sub>22</sub>), is expected to undergo biodegradation in the environment at a rate between that of the two structural end members of this series. Despite the slower rate of biodegradation implied by these study results, all of the fatty acids in the group are expected to undergo relatively rapid and ultimate biodegradation in the environment.

## Bioaccumulation

Chemicals in this group will release lead(2+) ions, which bioaccumulate in most organisms.

The lead(2+) ions released from these chemicals may pose a bioaccumulation hazard depending on the bioavailability of the metal ion under the prevailing exposure conditions. Conventional measures of bioaccumulation as applied to organic chemicals are not appropriate for metal ions. These measures do not consider the potential for metals to accumulate in specific tissues, the physiological mechanisms available to organisms to regulate internal metal concentrations, and the influence of environmental factors (US EPA, 2007). Nevertheless, lead and lead compounds are globally acknowledged to pose a significant bioaccumulation concern (UNEP, 2010; 2014).

The fatty acid components of the chemicals in this group are not expected to pose a bioaccumulation hazard. They are dietary fatty acids which are naturally ubiquitous in the environment and organisms, and their storage in body tissues is an essential biological process which is regulated by homeostatic controls (NRC, 2005).

## Transport

The chemicals in this group and their ionic components have a low potential for long-range transport.

The lead(2+) salts of fatty acids are ionic solids that do not partition to the atmospheric compartment, where long-range transport typically occurs. The fatty acid constituents of these salts are rapidly biodegradable and therefore not subject to long-range transport. The ionic lead components of these salts may have some limited mobility in water due to complexation by dissolved organic matter, but lead released into water and onto soil will eventually become bound to sediment or soil particles (UNEP, 2010).

## Predicted Environmental Concentration (PEC)

A PEC was not calculated for the chemicals or their ionic components addressed under this assessment.

Diffuse emissions of soluble lead(2+) from PVC articles can occur. Studies which have focused on the implications of this for drinking water supplies have highlighted the potential for lead to be released from PVC pipes to water (Al-Malack, 2001; Koh, et al., 1991; Lasheen, et al., 2008). However, other studies have concluded that extraction of lead from PVC pipes by running water rapidly declines shortly after initial use and that lead released by this process makes a relatively small contribution to total lead emissions to the environment (Smith, 1998).

The majority of lead distearate which is used industrially is expected to enter landfills incorporated into PVC articles disposed of at the end of their useful life. A number of studies have examined the release of lead from PVC articles in landfills and they conclude that most of the lead contained in PVC is expected to remain bound within the polymer matrix. Furthermore, a study conducted in the European Union found that the lead from lead compounds used in PVC pipes and cables represents approximately 3% of total lead content in municipal solid waste (ARGUS, et al., 2000). It is noted that this value may also include other lead-based compounds that are used in PVC but are not addressed under this assessment. A similar Australian review concluded that the amount of lead (in all forms) released to the environment due to use of lead compounds in PVC was relatively small compared to lead contributions from other sources (Smith, 1998).

## Environmental Effects

No ecotoxicity data specific to the chemicals addressed under this assessment are available. However, it is expected that the environmental effects of sparingly soluble lead(2+) salts of fatty acids will be determined principally by the release of lead(2+) ions into the environment. Lead is recognised to be a highly toxic environmental contaminant (UNEP, 2010; 2014). The effects assessment of the chemicals in this group has therefore primarily focused on the extensive body of information regarding the ecotoxicity of lead and inorganic lead compounds.

## Effects on Aquatic Life

Bioavailable forms of lead(2+) are very toxic to aquatic life in short and long term exposures.

## Toxicity of lead(2+)

The speciation of lead(2+) is strongly dependent on water chemistry, which can have a dramatic effect on the bioavailability and therefore toxicity of lead. For example, lead(2+) is more toxic to freshwater species in soft water than hard water. Toxicity is also higher in acidic freshwater when compared to alkaline freshwater. However, toxicity is lower to marine species than for freshwater species (ANZECC, 2000a). These factors and their effects should be noted when evaluating the toxicity of lead(2+) salts.

The following acute median lethal concentrations (LC50s) and median effective concentration (EC50) for model organisms across three trophic levels for lead (CAS RN 7439-92-1) were reported in the Registration Dossier for the chemical under the

European Union REACH legislation (ECHA, 2014f). All studies were conducted on freshwater species using soluble salts of lead(2+):

Taxon	Endpoint	Method
Fish	96 h LC50 = 108 µg Pb/L	Experimental <i>Pimephales promelas</i> (Fathead minnow) ASTM Method; Flow through CaCO <sub>3</sub> = 43.9 mg/L, pH = 7.4
Invertebrates	48 h LC50 = 73.6 µg Pb/L	Experimental <i>Ceriodaphnia dubia</i> (Water flea) US EPA Method; Semi-static CaCO <sub>3</sub> = 16.4 mg/L, pH = 5.7
Algae	48 h EC50 = 23.1 µg Pb/L	Experimental <i>Pseudokirchneriella subcapitata</i> (Green algae) OECD TG 201; Static CaCO <sub>3</sub> = 24 mg/L, pH = 7.3 Reduced growth rate observed

The chronic toxicity of lead(2+) to freshwater and marine species was critically evaluated for the compilation of water quality trigger values for environmental contaminants in the Australian and New Zealand Guidelines for Fresh and Marine Water Quality. Only studies that considered the complex relationship between water chemistry and toxicity were considered, and no-observed effect concentration (NOEC) equivalent values were derived (ANZECC, 2000a). The equivalent NOEC values available for sensitive freshwater fish and invertebrate species are reported below, along with an effective concentration for 10% of the test population (EC10) value for algae as reported in the Registration Dossier for lead under the European Union REACH legislation (ANZECC, 2000a; ECHA, 2014f):

Taxon	Endpoint	Method
Fish	NOEC = 5.65 µg Pb/L	Experimental <i>Lepidomeda vittatus</i> (Little Colorado spinedace) CaCO <sub>3</sub> = 30 mg/L

Taxon	Endpoint	Method
Invertebrates	NOEC = 19.5 µg Pb/L	Experimental <i>Daphnia magna</i> (Water flea) CaCO <sub>3</sub> = 30 mg/L
Algae	48 h EC10 = 4.5 µg Pb/L	Experimental <i>Pseudokirchneriella subcapitata</i> (Green algae) OECD TG 201; Static CaCO <sub>3</sub> = 24 mg/L, pH = 7.3 Reduced growth rate observed

## Toxicity of long-chain fatty acids

The long-chain fatty acid constituents of these chemicals have only moderate toxicity to aquatic life in short and long term exposures.

The following acute LC50 value and EC50 values for model freshwater organisms across three trophic levels for (a) oleic acid (CAS RN 112-80-1) and (b) docosanoic acid (CAS RN 112-85-6) were reported in the databases included in the OECD QSAR Toolbox (LMC, 2013). Additional available toxicity values similarly reflect moderate to low ecotoxicity for these long-chain fatty acids (LMC, 2013; NITE, 2014):

Taxon	Endpoint	Method
Fish	(a) 96 h LC50 = 205 mg/L	Experimental <i>Pimephales promelas</i> (Fathead minnow)
Invertebrates	(b) 48 h EC50 = 5 mg/L	Experimental <i>Daphnia magna</i> (Water flea) OECD TG 202 Immobilisation observed
Algae	(b) 72 h EC50 = 5 mg/L	Experimental <i>Pseudokirchneriella subcapitata</i> (Green algae) Reduced growth rate observed

The following chronic NOEC values for model organisms across two trophic levels for (a) oleic acid, (b) docosanoic acid and (c) palmitic acid (CAS RN 57-10-3) were reported in the databases included in the OECD QSAR Toolbox (LMC, 2013). Additional available threshold toxicity values suggest a similar degree of chronic toxicity to aquatic invertebrates and algae is not a group-wide trend, with comparable NOEC values greater than 100 mg/L available for 12-hydroxystearic acid (CAS RN 106-14-9) (LMC, 2013; NITE, 2014):

Taxon	Endpoint	Method
Invertebrates	(a) 21 d NOEC = 0.32 mg/L	Experimental <i>Daphnia magna</i> (Water flea) OECD TG 211 Reproductive toxicity observed
Algae	(a) 72 h NOEC = 2.6 mg/L (b) 72 h NOEC = 4.1 mg/L (c) 72 h NOEC = 0.6 mg/L	Experimental <i>Pseudokirchneriella subcapitata</i> (Green algae) OECD TG 201 Reduced growth rate observed

## Effects on Sediment-Dwelling Life

Bioavailable forms of lead(2+) are very toxic to sediment-dwelling organisms.

### Toxicity of lead(2+)

The toxicity of lead to sediment-dwelling organisms has been summarised in the rationale and background information to the Australian and New Zealand Guidelines for Fresh and Marine Water Quality. Exposure to sediment contaminants for benthic organisms is assumed to principally involve pore waters (ANZECC, 2000b). The available NOEC value equivalents for select sediment-dwelling organisms (corrected to a CaCO<sub>3</sub> concentration of 30 mg/L) include a NOEC of 5.1 micrograms of lead per litre (µg Pb/L) for the amphipod *Gammarus pseudolimnaeus* and a NOEC of 28 µg Pb/L for the midge *Tanytarsus dissimilis* (ANZECC, 2000a). However, toxicity is highly dependent on sediment chemistry, which is complex and depends on a range of variable environmental parameters such as dissolved oxygen, pH and the geochemistry of sediment particles (ANZECC, 2000b). In particular, the presence of natural organic matter and the formation of sulfides will affect lead bioavailability. These factors must also be considered when determining the toxicity of lead(2+) to sediment-dwelling species.

### Toxicity of long-chain fatty acids

No suitable data were available to evaluate the effects of the fatty acid constituents of these chemicals on sediment-dwelling organisms. However, the release of lead(2+) ions to the sediment compartment is expected to dominate ecotoxicity concerns for these salts.

## Effects on Terrestrial Life

Bioavailable forms of lead(2+) are toxic to terrestrial organisms.

## Toxicity of lead(2+)

The toxicity of lead to terrestrial organisms has been summarised in Schedule B5c of the *National Environment Protection (Assessment of Site Contamination) Measure 1999* (Cwlth). Lead is strongly adsorbed to organic matter in soil and therefore bioavailability is typically limited. Mean NOEC values for terrestrial invertebrates range from 129 milligrams of lead per kilogram of soil (mg Pb/kg soil) for the earthworm *Dendrobaena rubida* to 1797 mg Pb/kg soil for the springtail *Fautrix candida*. Plants are also susceptible to lead toxicity, with mean NOEC values in a range from 50 mg Pb/kg soil for barley (*Hordeum vulgare*) to 546 mg Pb/kg soil for the Loblolly pine (*Pinus taeda*). Reproductive toxicity and impaired growth are listed as the most common toxic effects (Commonwealth of Australia, 1999).

## Toxicity of long-chain fatty acids

No suitable data were available to evaluate the effects of the fatty acid constituents of these chemicals on terrestrial organisms. However, release of lead(2+) ions to the terrestrial compartment is again expected to dominate ecotoxicity concerns for these salts.

## Predicted No-Effect Concentration (PNEC)

The primary environmental effects of the chemicals in this group are expected to be caused by the release of lead(2+) ions.

In place of the PNECs for aquatic and sediment compartments for this ionic component, the trigger values published for lead in the Australian and New Zealand Guidelines for Fresh and Marine Water Quality have been used. These values represent thresholds above which further assessment of potential toxicity may be required to ensure environmental quality. Although they were not developed for risk assessment purposes, they nevertheless provide important reference values for evaluating the risks posed by anthropogenic emissions of lead to the Australian environment. For freshwater ecosystems with low water hardness (30 mg/L CaCO<sub>3</sub>), a high reliability trigger value for protection of 95% of species has been determined to be 3.4 µg Pb/L. The equivalent value for protection of marine species is 4.4 µg Pb/L. For the sediment compartment, the trigger value is 50 mg Pb/kg, dry weight (ANZECC, 2000c).

In place of a PNEC for the soil compartment for lead(2+), the added contaminant limit published for lead in the *National Environment Protection (Assessment of Site Contamination) Measure 1999* (Cwlth) has been used. This value represents the level above which further investigation or evaluation is required after considering naturally occurring background levels. For soil where the contaminant has been present for less than two years, the contaminant limit for protection of 99% of species is 110 mg added Pb/kg soil (Commonwealth of Australia, 1999).

## Categorisation of Environmental Hazard

It is not currently possible to categorise the environmental hazards of metals and other inorganic chemicals according to standard persistence, bioaccumulation and toxicity (PBT) hazard criteria. These criteria were developed for organic chemicals and do not take into account the unique properties of inorganic substances and their behaviour in the environment and biota (UNECE, 2007; US EPA, 2007). Therefore, environmental hazard categorisation according to domestic PBT criteria has only been performed for the organic components of the chemicals in the 1:2 Lead(2+) Salts of Long-Chain Carboxylic Acids group, as presented below:

### Persistence

Not Persistent (Not P). Based on results obtained from various biodegradation studies which indicate rapid and ultimate biodegradation of long-chain fatty acids, the organic components of all chemicals in this group are categorised as Not Persistent.

### Bioaccumulation

Not Bioaccumulative (Not B). Based on the essential biological functions of long-chain fatty acids as dietary fats and the associated homeostatic controls, the organic components of all chemicals in this group are categorised as Not Bioaccumulative.

## Toxicity

Not Toxic (Not T). Based on available acute ecotoxicity values greater than 1 mg/L, and chronic ecotoxicity values greater than 0.1 mg/L, the organic components of all chemicals in this group are categorised as Not Toxic.

## Summary

The organic components of octadecanoic acid, lead(2+) salt; octadecanoic acid, 12-hydroxy-, lead(2+) salt (2:1); hexadecanoic acid, lead(2+) salt; eicosanoic acid, lead(2+) salt (2:1); docosanoic acid, lead(2+) salt; 9-hexadecenoic acid, lead(2+) salt, (Z)-; 9-octadecenoic acid, 12-hydroxy-, lead(2+) salt (2:1), [R-(Z)]; 9-octadecenoic acid, lead(2+) salt, (Z)-; and 9,12-octadecadienoic acid, (Z,Z)-, lead(2+) salt are all categorised as:

- Not P
- Not B
- Not T

An environmental hazard categorisation for the inorganic lead(2+) component of the chemicals in this group has not been assigned for the reasons discussed above. Nevertheless, there is an international consensus that bioavailable forms of lead are highly toxic to the environment and that lead bioaccumulates in most organisms (UNEP, 2010; 2014).

## Risk Characterisation

Risk quotients (RQs) have not been calculated for these chemicals.

The chemicals in this group contain lead(2+) ions, which can be released into the environment from industrial applications. Anthropogenic emissions of lead to the environment are acknowledged to be of concern, both domestically and internationally, due to the bioaccumulation properties and high toxicity of bioavailable forms of lead. Environmental contamination by lead is of particular concern because high exposure to lead compounds can cause neurological defects and reproductive toxicity across a range of organisms, including humans, plants and animals (UNEP, 2010; 2014).

However, the release of lead to the environment in Australia from industrial uses of chemicals in this group is expected to be limited in volume and diffuse in nature. Available research suggests that the contribution of lead from lead salts of long-chain fatty acids to total environmental lead loads is comparatively small. Furthermore, recent changes in the industrial manufacture of PVC resins have dramatically reduced the consumption of these lead salts for their main industrial use as stabilisers. The risks from lead released from PVC articles is expected to decline further as lead salts of long-chain fatty acids are finally phased out of use as stabilisers.

## Key Findings

The main industrial use of chemicals in the 1:2 Lead(2+) Salts of Long-Chain Carboxylic Acids group is expected to be as stabilisers in the manufacture of PVC articles. The most important member of this group is lead distearate which was previously used in significant volumes as a stabiliser both domestically and internationally. However, available information from domestic and international sources indicates that the use of lead(2+) salts of long-chain fatty acids as stabilisers for PVC has largely been phased out as a result of industry initiatives.

The principal environmental concern for industrial uses of lead(2+) salts of long-chain fatty acids is the potential for release of soluble forms of ionic lead from manufactured PVC articles. This poses a concern because lead and lead compounds are considered to be highly significant environmental contaminants. However, an analysis of the risks arising from the industrial use

of these chemicals indicates that the quantity of lead released from PVC articles containing lead-based stabilisers makes a relatively small contribution to total anthropogenic lead emissions.

The chemicals in this group are considered to pose a low and declining risk to the environment based on the limited contribution to total anthropogenic lead emissions from lead-based stabilisers, and the phase out of use of these chemicals as stabilisers in PVC. No further assessment of the environmental risks of these chemicals is currently required.

The organic components of the chemicals in the 1:2 Lead(2+) Salts of Long-Chain Carboxylic Acids group are not PBT substances according to domestic environmental hazard criteria.

## Recommendations

The nine chemicals in the 1:2 Lead(2+) Salts of Long-Chain Carboxylic Acids group are not prioritised for further assessment under the IMAP framework.

## Environmental Hazard Classification

In addition to the categorisation of environmental hazards according to domestic environmental thresholds presented above, the classification of the environmental hazards of octadecanoic acid, lead(2+) salt; octadecanoic acid, 12-hydroxy-, lead(2+) salt (2:1); hexadecanoic acid, lead(2+) salt; eicosanoic acid, lead(2+) salt (2:1); docosanoic acid, lead(2+) salt; 9-hexadecenoic acid, lead(2+) salt, (Z)-; 9-octadecenoic acid, 12-hydroxy-, lead(2+) salt (2:1), [R-(Z)]; 9-octadecenoic acid, lead(2+) salt, (Z)- and 9,12-octadecadienoic acid, (Z,Z)-, lead(2+) salt according to the third edition of the United Nations' Globally Harmonised System of Classification and Labelling of Chemicals (GHS) is presented below (UNECE, 2009):

Hazard	GHS Classification (Code)	Hazard Statement
Acute Aquatic	Category 1 (H400)	Very toxic to aquatic life
Chronic Aquatic	Category 1 (H410)	Very toxic to aquatic life with long lasting effects

The classification of octadecanoic acid, lead(2+) salt (lead distearate) was conducted based on the data available from a study conducted in accordance with the OECD Transformation and Dissolution Protocol and the ecotoxicological data identified in this assessment (ECHA, 2014d; 2014f; LMC, 2013). The chemical is classified as Acute Aquatic Category 1 as the dissolved metal concentration obtained from the lowest loading level after seven days (27.2 µg Pb/L) exceeds the acute toxicity value (48 h EC50) available for the algae *Pseudokirchneriella subcapitata*. Similarly, the dissolved metal concentration obtained from the lowest loading level after 28 days (41.2 µg Pb/L) exceeds all chronic toxicity endpoints listed above for aquatic species. Also, lead(2+) bioaccumulates in aquatic organisms. Hence, this salt is categorised as Chronic Aquatic Category 1 according to the GHS classification strategy for metals and metal compounds (UNECE, 2007).

Based on the available  $K_{sp}$  for hexadecanoic acid, lead(2+) salt (lead dipalmitate), this chemical is considered more soluble than lead distearate. Therefore, this chemical will be considered more toxic than lead distearate under the GHS. However, lead distearate has been classified under the highest environmental hazard categories available. Therefore, this more soluble chemical has been classified under the same aquatic hazard categories as lead distearate.

There are insufficient solubility data available for the remaining chemicals in this group to classify their aquatic toxicity hazards according to the procedure outlined above. In accordance with the classification guidance provided for metals and metal compounds in Annex 9 of the GHS, these chemicals are assumed to be sufficiently soluble to release lead(2+) ions at concentrations which exceed the relevant ecotoxicological endpoints (UNECE, 2007). Therefore, these remaining chemicals are all classified as Acute Aquatic Category 1 and Chronic Aquatic Category 1 based on the available toxicity data for lead.

It is preferable to classify the hazard posed by metals and metal compounds using the findings of a study conducted in accordance with the OECD Transformation and Dissolution protocol (UNECE, 2007). Therefore, should a study conducted in accordance with this protocol suggest a lower hazard classification is warranted, these chemicals may be reclassified as appropriate.

## References

- Al-Malack MH (2001). Migration of lead from unplasticized polyvinyl chloride pipes. *Journal of Hazardous Materials*, **B82**, pp 263.
- ANZECC (2000a). 8.3 Toxicants. In: Australian and New Zealand Environment and Conservation Council, *Australian and New Zealand guidelines for fresh and marine water quality: Volume 2 - Aquatic ecosystems - rationale and background information*. Canberra, Australia. Available at <http://www.environment.gov.au>.
- ANZECC (2000b). 8.4 Sediment Quality Guidelines. In: Australian and New Zealand Environment and Conservation Council, *Australian and New Zealand guidelines for fresh and marine water quality: Volume 2 - Aquatic ecosystems - rationale and background information*. Canberra, Australia. Available at <http://www.environment.gov.au>.
- ANZECC (2000c). *Australian and New Zealand Guidelines for Fresh and Marine Water Quality*. Australian and New Zealand Environment and Conservation Council, Sydney, Australia. <http://www.environment.gov.au>.
- ARGUS, et al. (2000). *The Behaviour of PVC in Landfill*. European Commission, Brussels, Belgium. <http://ec.europa.eu>.
- Australian Government Department of the Environment (2013a). *Understanding NPI data*. Australian Government Department of the Environment, Canberra, Australia. Accessed 11 March 2014 at <http://www.npi.gov.au>.
- Australian Government Department of the Environment (2013b). *NPI substance list and thresholds*. Australian Government Department of the Environment, Canberra, Australia. Accessed 11 March 2014 at <http://www.npi.gov.au>.
- Calhoun SF (1960). Antiwear and Extreme Pressure Additives for Grease. *Annual Meeting of the American Society of Lubrication Engineers*, Ohio, USA. Accessed at <http://www.stle.org>
- Chemiplas (2012). *Submission to the Anti-Dumping Commission in relation to ITR 184*. Australian Customs and Border Protection Services, Canberra, Australia. <http://www.adcommission.gov.au>.
- Commonwealth of Australia (1999). *National Environment Protection (Assessment of Site Contamination) Measure 1999*.
- Commonwealth of Australia (2013). *Poisons Standard 2013*. Poisons Standard. Enacted on 23 August 2013.
- ECHA (2013). *Candidate List of Substances of Very High Concern for Authorisation*. European Chemicals Agency, Helsinki, Finland. Accessed 17 March 2014 at <http://echa.europa.eu>.
- ECHA (2014a). *Pre-registered substances*. European Chemicals Agency, Helsinki, Finland. Accessed 1 May 2014 at <http://echa.europa.eu>.
- ECHA (2014b). *Registered Substances*. European Chemicals Agency, Helsinki, Finland. Accessed 9 May 2014 at <http://echa.europa.eu>.
- ECHA (2014c). *Substances to be registered*. European Chemicals Agency, Helsinki, Finland. Accessed 1 May 2014 at <http://echa.europa.eu>.
- ECHA (2014d). *Fatty acids, C16-18, lead salts*. European Chemicals Agency, Helsinki, Finland. Accessed 26 May 2014 at <http://apps.echa.europa.eu>.
- ECHA (2014e). *Dioxobis(stearato)trilead*. European Chemicals Agency, Helsinki, Finland. Accessed 26 May 2014 at <http://apps.echa.europa.eu>.
- ECHA (2014f). *Lead*. European Chemicals Agency, Helsinki, Finland. Accessed 29 May 2014 at <http://apps.echa.europa.eu>.

Environment Canada (2013a). *Status of Prioritized Substances*. Environment Canada, Gatineau, Canada. Accessed 1 May 2014 at <http://www.ec.gc.ca>.

Environment Canada (2013b). *Search Engine for Chemicals and Polymers*. Environment Canada, Gatineau, Canada. Accessed 1 May 2014 at <http://www.ec.gc.ca>.

Environment Canada (2013c). *Search Engine for the Results of DSL Categorisation*. Environment Canada, Gatineau, Canada. Accessed 21 November 2013 <http://www.ec.gc.ca>.

EPT (2003). *End-of-life Environmental Issues with PVC in Australia*. ExcelPlas Polymer Technology, Melbourne, Australia. <http://www.environment.gov.au>.

ESPA (2014). *Stabilisers - What's new?* European Stabiliser Producers Association, Brussels, Belgium. <http://www.stabilisers.eu>.

Government of Canada (2005). *Surface Coatings Materials Regulations*. Enacted on 19 April 2005.

Health Canada (2013). *Final Human Health State of the Science Report on Lead*. Health Canada, Ottawa, Canada. <http://www.hc-sc.gc.ca>.

Hunter KA and Liss PS (1976). Measurement of the Solubility Products of Various Metal Ion Carboxylates. *Journal of Electroanalytical Chemistry*, **73**, pp 347.

Keml (2014). *Keml-stat*. Swedish Chemicals Agency (Swedish: Kemikalieinspektionen), Sundbyberg, Sweden. Accessed 1 May 2014 at <http://apps.kemi.se/kemistat/start.aspx?sprak=e>

Koh L, et al. (1991). Factors Affecting the Leaching of Lead from UPVC Pipes. *Fourth Symposium on our Environment*, Singapore. Accessed at <http://link.springer.com>

Lasheen M, et al. (2008). Factors influencing lead and iron release from some Egyptian drinking water pipes. *Journal of Hazardous Materials*, **160**, pp 675.

LMC (2013). *The OECD QSAR Toolbox for Grouping Chemicals into Categories*, v 3.1. Laboratory of Mathematical Chemistry, University "Prof. Dr. Assen Zlatarov", Burgas, Bulgaria. Available at <http://oasis-lmc.org>.

Mauchauffee S, et al. (2008). Determination of the Solubility Products in Water at 20°C of 32 Metallic Carboxylates. *Industrial and Engineering Chemistry Research*, **47**, pp 7533.

NICNAS (2006). *Australian High Volume Industrial Chemical List*. National Industrial Chemical Notification and Assessment Scheme, Sydney, Australia. <http://www.nicnas.gov.au>.

NICNAS (2007). *Lead Compounds in Industrial Surface Coatings and Inks*. National Industrial Chemical Notification and Assessment Scheme, Sydney, Australia. <http://www.nicnas.gov.au>.

NICNAS (2013). *Human Health Tier II Assessment for Lead Salts of Selected Fatty Acids*. National Industrial Chemicals Notification and Assessment Scheme, Sydney, Australia. Accessed 2 May 2014 at <http://www.nicnas.gov.au>.

NICNAS (2014). *AICS Listing: CAS Number 1072-35-1*. National Industrial Chemical Notification and Assessment Scheme, Sydney, Australia. Accessed 12 May 2014 at <http://www.nicnas.gov.au>.

NITE (2014). *Japan Chemicals Collaborative Knowledge Database*. National Institute of Technology and Evaluation, Tokyo, Japan. Accessed 19 March 2014 at <http://www.safe.nite.go.jp>.

Nora A and Koenen G (2012). Metallic Soaps. In: Bellussi G, et al., *Ullmann's Encyclopedia of Industrial Chemistry*. Wiley. Available at <http://onlinelibrary.wiley.com>.

Nordic Council of Ministers (2013). *Substances in Preparations in Nordic Countries (SPIN)*. Chemical Group, Nordic Council of Ministers, Copenhagen, Denmark. Downloaded 9 November 2012. Available at <http://www.spin2000.net>.

NRC (2005). Dietary Fats: Total Fat and Fatty Acids. In: National Research Council, *Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein and Amino Acids (Macronutrients)*. The National Academies Press,

Washington DC, USA. Available at <http://www.nap.edu>.

OECD (2004). *Emission Scenario Document on Lubricants and Lubricant Additives*. Paris, France. <http://search.oecd.org>.

OECD (2012). *Assessment of Chemicals: Manual for the Assessment of Chemicals, Chapter 1*. Organisation for Economic Cooperation and Development, Paris, France. Accessed October, 2013 <http://www.oecd.org>.

OECD (2013). *OECD Existing Chemicals Database*. Organisation for Economic Cooperation and Development, Paris, France. Accessed 13 November 2013 <http://webnet.oecd.org>.

PVC Europe (2014). *Lead Stabilisers*. PVC Europe, Brussels, Belgium. Accessed 9 May 2014 at <http://www.pvc.org>.

Schwartz M (2012). *Encyclopedia of Materials, Parts and Finishes*. CRC Press, Florida, USA.

Shugarman A (2003). *Monitoring Active Sulfur in EP Gear Oils - And Other Options for Monitoring EP Additive Depletion*. Noria Corporation, Tulsa, USA. Accessed at <http://www.machinerylubrication.com>.

Smith R (1998). *The Environmental Aspects of the Use of PVC in Building Products*. CSIRO Publishing, Melbourne, Australia.

Tilley K (2013). *Australian PVC industry phases out lead*. Plastics News. Accessed 5 May 2014 at <http://www.plasticsnews.com>.

UNECE (2007). *Annex 9: Guidance on Hazards to the Aquatic Environment*. United Nations Economic Commission for Europe, Geneva, Switzerland. <http://www.unece.org>.

UNECE (2009). *Globally Harmonised System of Classification and Labelling of Chemicals (GHS), 3rd Revised Edition*. United Nations Economic Commission for Europe, Geneva, Switzerland. Accessed 12 November 2013 at <http://www.unece.org>

UNEP (1985). *The Vienna Convention for the Protection of the Ozone Layer and its Montreal Protocol on Substances that Deplete the Ozone Layer*. United Nations Environment Programme, Ozone Secretariat, Nairobi, Kenya. Accessed 12 November 2013 at <http://ozone.unep.org>.

UNEP (2001). *The Stockholm Convention on Persistent Organic Pollutants*. United Nations Environment Programme, Secretariat of the Stockholm Convention, Châtelaine, Switzerland. Accessed 12 November 2013 at <http://www.pops.int>.

UNEP (2010). *Final review of scientific information on lead*. United Nations Environment Programme, Nairobi, Kenya. <http://www.unep.org>.

UNEP (2011). *Operational Framework for the Global Alliance to Eliminate Lead Paint*. United Nations Environment Programme, Nairobi, Kenya. <http://www.unep.org>.

UNEP (2014). *Lead and Cadmium*. United Nations Environment Programme, Nairobi, Kenya. Accessed 5 March 2014 at <http://www.unep.org>.

UNEP & FAO (1998). *The Rotterdam Convention on the Prior Informed Consent procedure for Certain Hazardous Chemicals and Pesticides in International Trade*. United Nations Environment Programme and Food and Agriculture Organization of the United Nations, Châtelaine, Switzerland. Accessed 12 November 2013 at <http://www.pic.int>.

United States Government (2008). *Consumer Product Safety Improvement Act of 2008*. CPSIA. Enacted on 14 August 2008.

US EPA (1999). *Understanding Variation in Partition Coefficient, K<sub>d</sub>, Values - Volume II: Review of Geochemistry and Available K<sub>d</sub> Values for Cadmium, Cesium, Chromium, Lead, Plutonium, Radon, Strontium, Thorium, Tritium (3H), and Uranium*. United States Environmental Protection Agency, Washington DC, USA. <http://www.epa.gov>.

US EPA (2007). *Framework for Metals Risk Assessment*. United States Environmental Protection Agency, Washington DC, USA. <http://www.epa.gov>.

US EPA (2012). *Is My Facility's Six-Digit NAICS Code a TRI-Covered Industry?* United States Environmental Protection Agency, Washington DC, USA. Accessed 17 April 2013 at <http://www2.epa.gov>.

US EPA (2014a). *Basics of TRI Reporting*. United States Environmental Protection Agency, Washington DC, USA. Accessed 3 April 2014 at <http://www2.epa.gov>.

US EPA (2014b). *TRI-Listed Chemicals*. United States Environmental Protection Agency, Washington DC, USA. Accessed 3 April 2014 at <http://www2.epa.gov>.

Vinyl 2010 (2006). *Vinyl 2010: The Voluntary Commitment of the PVC Industry*. Vinyl 2010, Brussels, Belgium. <http://www.vinylplus.eu>.

Vinyl 2010 (2011). *Vinyl 2010: 10 Years*. Vinyl 2010, Brussels, Belgium. <http://www.vinylplus.eu>.

Vinyl Council Australia (2012). *PVC Product Stewardship Program*. Vinyl Council Australia, Melbourne, Australia. <http://vinyl.org.au>.

Vinyl Council Australia (2014). *Signatories*. Vinyl Council Australia, Melbourne, Australia. Accessed 26 May 2014 at <http://www.vinyl.org.au/pvc-the-environment/product-stewardship/list-of-signatories>

*Last update 4 July 2014*

Share this page