



**Australian Government**

**Department of Health**


Australian Industrial Chemicals Introduction Scheme

# **Galaxolide and a related polycyclic musk**

## **Evaluation statement**

**28 May 2021**

**Draft**



# Table of contents

## Contents

AICIS evaluation statement .....	4
Subject of the evaluation.....	4
Chemicals in this evaluation.....	4
Reason for the evaluation .....	4
Parameters of evaluation .....	4
Summary of evaluation .....	4
Summary of introduction, use and end use.....	4
Environment.....	5
Conclusions .....	6
Recommendations .....	6
Environment.....	6
Supporting information .....	7
Rationale .....	7
Chemical identity .....	7
Relevant physical and chemical properties .....	9
Introduction and use .....	10
Australia .....	10
International .....	10
Existing Australian regulatory controls .....	11
Environment.....	11
International regulatory status.....	11
United Nations.....	11
United States of America.....	12
Environmental exposure .....	12

Environmental fate .....	12
Predicted environmental concentration (PEC) .....	15
Environmental effects .....	17
Effects on aquatic life .....	17
Effects on sediment dwelling life.....	18
Effects on terrestrial life.....	19
Endocrine effects/activity.....	19
Predicted no-effect concentration (PNEC).....	20
Categorisation of environmental hazard.....	20
Persistence .....	20
Bioaccumulation .....	20
Toxicity.....	20
Environmental risk characterisation .....	20
Uncertainty.....	21
References .....	22

# AICIS evaluation statement

## Subject of the evaluation

Galaxolide and a related polycyclic musk.

## Chemicals in this evaluation

Name	CAS Registry Number
Cyclopenta[g]-2-benzopyran, 1,3,4,6,7,8-hexahydro-4,6,6,7,8,8-hexamethyl-	1222-05-5
1H-Naphtho[2,3-c]pyran, 3,4,6,7,8,9-hexahydro-4,6,6,9,9-pentamethyl-	1922-67-4

## Reason for the evaluation

The Evaluation Selection Analysis indicated a potential risk to the environment.

## Parameters of evaluation

This evaluation considers the environmental risks associated with the industrial uses of galaxolide (CAS RN 1222-05-5) and musk 89 (CAS RN 1922-67-4), two synthetic polycyclic musk fragrances. These chemicals have been assessed for their risks to the environment according to the following parameters:

- Default domestic introduction volumes of 100 tonnes per annum
- Industrial uses listed in the 'Summary of Use' section
- Expected emission into sewage treatment plants (STPs) due to consumer and commercial use.

These chemicals have been assessed as a group as they are structurally very similar and have similar use patterns.

## Summary of evaluation

### Summary of introduction, use and end use

Galaxolide is a common fragrance ingredient in a variety of cosmetic and consumer-use products worldwide, with a global use volume in the thousands of tonnes per year. There are no specific domestic introduction volume data available for galaxolide.

Galaxolide is used in the following products according to reported domestic and international use data:

- Personal care products
- Air care products
- Laundry and dishwashing products

- Cleaning and furniture care products
- Automotive products.

No domestic or international use or introduction information was identified for musk 89. It does not appear to be in widespread industrial use.

## Environment

### Summary of environmental hazards

According to domestic environmental hazard thresholds and based on the available data galaxolide and musk 89 are:

- Persistent (P)
- Not Bioaccumulative (not B)
- Toxic (T)

### Environmental hazard classification

The chemical satisfies the criteria for classification according to the Globally Harmonized System of Classification and Labelling of Chemicals (GHS) for environmental hazards as follows. This does not consider classification of physical hazards and health hazards.

Environmental Hazard	Hazard Category	Hazard Statement
Acute Aquatic	Acute Aq. – Cat. 1	H400: Very toxic to aquatic life
Chronic Aquatic	Chronic Aq. – Cat. 1	H410: Very toxic to aquatic life with long-lasting effects

### Summary of environmental risk

Galaxolide has global use volumes in the thousands of tonnes per year. It is used widely as a fragrance ingredient in personal care and other domestic use products and is released to wastewater as a normal part of its use pattern.

Galaxolide is toxic, has moderate bioaccumulation potential, and is categorised as persistent based on its long half-life in some soils and its degradation to persistent degradants. Primary degradation of galaxolide into galaxolidone (CAS RN 507442-49-1) and other polar degradants is rapid in a variety of environmental compartments. However, ultimate degradation of galaxolide has not been observed under environmentally relevant conditions.

A detailed analysis of the hazard characteristics of metabolites of galaxolide has not been conducted as part of this evaluation due to lack of data. However, all identified metabolites are more polar than galaxolide, and appear to be excreted rapidly by fish. These metabolites are therefore expected to have lower bioaccumulation potential than galaxolide, and likely do not have PBT characteristics.

Based on measured domestic and international concentrations in sediment, STP effluent and biosolids, galaxolide is expected to be present in Australian river, sediment and soil compartments at concentrations below the level of concern.

Musk 89 is a close structural analogue of galaxolide and is expected to have similar hazard characteristics. It is not expected to be present in the Australian environment at concentrations approaching the level of concern based on its apparent lack of widespread industrial use.

## Conclusions

The conclusions of this evaluation are based on the information described in the statement. Obligations to report additional information about hazards under section 100 of the *Industrial Chemicals Act 2019* apply.

The Executive Director is satisfied that the identified environment risks can be managed within existing risk management frameworks. This is provided that all requirements are met under environmental, workplace health and safety and poisons legislation as adopted by the relevant state or territory.

Although the chemicals are present at concentrations below the level of concern in the Australian environment, given the hazard characteristics of the chemicals there is a risk of adverse effects on the environment if levels increase. The proposed means of managing the risks identified during this evaluation are set out in the **Recommendations** section.

## Recommendations

### Environment

#### Industrial Chemicals Environmental Management Standard (IChEMS)

Schedule the chemicals under the *Industrial Chemicals Environmental Management (Register) Act 2021*.

# Supporting information

## Rationale

This evaluation considers environmental risks associated with the industrial uses of galaxolide and musk 89, two closely related synthetic tricyclic musks. The evaluation of these substances has been conducted as a group because they have known or potential applications as synthetic musk fragrances.

Galaxolide and musk 89 were developed as synthetic alternatives to natural musk chemicals (Heeringa and Beets, 1963). Galaxolide is a high production volume chemical internationally and is commonly found in personal care and household products (Reiner and Kannan, 2006). Its use in these products has potential to result in environmental exposure through emission to sewers following their use, followed by release to the environment in the treated effluents and biosolids produced by sewage treatment plants (STPs).

The Evaluation Selection Analysis (ESA) of galaxolide highlighted potential persistence, bioaccumulation, and toxicity (PBT) hazard characteristics, which indicate a high concern for the environment. This evaluation includes further refinement of the risk characterisation, and a more in-depth assessment of the available environmental hazard and exposure information for both galaxolide and its structural analogue, musk 89.

Environmental risks resulting from the use of other polycyclic musks as fragrance ingredients in Australia have previously been assessed under the Inventory Multi-tiered Assessment and Prioritisation (IMAP) framework established by the National Industrial Chemicals Notification and Assessment Scheme (NICNAS). Environment Tier II assessments are available for [Tonalide and Related Polycyclic Musks](#) (NICNAS, 2016) and [Celestolide and Related Polycyclic Musks](#) (NICNAS, 2017).

## Chemical identity

The two chemicals in this assessment are structural isomers. Galaxolide and musk 89 are isochroman derivatives fused with an alkyl-substituted ring; galaxolide has a pentamethyl-substituted cyclopentyl ring and musk 89 has a tetramethyl-substituted cyclohexyl ring.


The structure of galaxolide contains two chiral centres. However, the stereochemistry of this substance is not defined in its Australian Inventory of Industrial Chemicals (Inventory) name. Technical galaxolide contains a mixture of the four stereoisomers in approximately equal proportions (Wang, et al., 2013).


The structure of musk 89 contains a single chiral centre, with its stereochemistry similarly undefined in its Inventory name. While no information was identified, it is assumed that technical mixtures of musk 89 are composed of a racemic (equal) mixture of two enantiomers.

Purified galaxolide (mixture of stereoisomers) is a crystalline solid whereas the technical mixture is a viscous liquid. The technical mixture has a purity of 70–80%, and is sold under the same name and CAS RN as its major component. The remaining 20–30% comprises four structural isomers of galaxolide that are by-products of the synthetic procedure. A typical mixture contains 6–16% combined of two isomers that each contain an ethyl group on the cyclopentyl ring in one of two possible positions (CAS RNs 78448-48-3 and 78448-49-4), and

4–16% combined of two isomers that differ from galaxolide in the position of the fused cyclopentyl ring (CAS RNs 114109-63-6 and 114109-62-5) (ECB, 2008; HERA, 2004; Sprecker, 1987). None of these structural isomers are listed on the Inventory. Musk 89 is synthesised via a distinct synthetic procedure and is not an impurity in the galaxolide technical mixture.

Due to difficulties using the viscous technical galaxolide in formulating fragrance blends, galaxolide is available commercially as a solution containing 65 parts by weight of the technical mixture and 35 parts by weight of solvent. Common solvents include diethyl phthalate (DEP, CAS RN 84-66-2), benzyl benzoate (BB, CAS RN 120-51-4) and isopropyl myristate (IPM, CAS RN 110-27-0). The resulting solutions are referred to as galaxolide 50 DEP, BB or IPM, respectively, with the '50' referring to the percentage of pure galaxolide (ECB, 2008).

CAS RN	1222-05-5
Chemical name	cyclopenta[ <i>g</i> ]-2-benzopyran, 1,3,4,6,7,8-hexahydro-4,6,6,7,8,8-hexamethyl-
Synonyms	galaxolide HHCB abbalide pearlide chromanolide hexamethylindanopyran
Structural formula	
Molecular formula	C <sub>18</sub> H <sub>26</sub> O
Molecular weight (g/mol)	258.4
SMILES	C1(C)(C)c2c(C(C)(C)C1C)cc1c(C(C)COC1)c2
Chemical description	-

CAS RN	1922-67-4
Chemical name	1 <i>H</i> -naphtho[2,3- <i>c</i> ]pyran, 3,4,6,7,8,9-hexahydro-4,6,6,9,9-pentamethyl-
Synonyms	musk 89
Structural formula	
Molecular formula	C <sub>18</sub> H <sub>26</sub> O
Molecular weight (g/mol)	258.4
SMILES	<chem>C1=C3C(=CC2=C1COCC2C)C(CCC3(C)C)(C)C</chem>
Chemical description	-

## Relevant physical and chemical properties

Measured physical and chemical property data for galaxolide were retrieved from the registration dossier for the chemical submitted under the Registration, Evaluation, Authorisation and Restriction of Chemicals (REACH) legislation in the European Union (EU) (REACH, 2011), the EU Risk Assessment Report (EU RAR) for galaxolide (ECB, 2008) and the patent literature (Heeringa and Beets, 1963).

There are few measured values available for the physical and chemical properties of musk 89. The melting point was retrieved from the patent literature (Heeringa and Beets, 1963), the values for other chemical properties were calculated using standard quantitative structure-property relationships (QSPR), and the Henry's Law constant was calculated from measured values for water solubility and vapour pressure (US EPA, 2017a):

Chemical	Galaxolide	Musk 89
Physical form	Solid	Solid
Melting point	53.0–53.5°C (exp.)	44.7–45.4°C (exp.)
Boiling point	318°C (exp.)	330°C (calc.)
Vapour pressure	0.0727 Pa (exp.)	0.0341 Pa (calc.)
Water solubility	1.75 mg/L (exp.)	0.11 mg/L (calc.)
Henry's Law constant	36.9 Pa·m <sup>3</sup> /mol (exp.)	80.2 Pa·m <sup>3</sup> /mol (calc.)
Ionisable in the environment?	No	No
pK <sub>a</sub>	N/A	N/A
log K <sub>ow</sub>	5.9 (exp.)	6.3 (calc.)

Information is not available on whether the test substance used for measurements of physical and chemical properties was pure crystalline galaxolide, or the liquid technical mixture. It is likely that melting point measurements were made on the purified material, since the technical mixture is a liquid at room temperature. For all other measured data, including standard tests on environmental hazard characteristics, the precise nature of the galaxolide test material is unknown and is a source of uncertainty in this evaluation.

Since musk 89 is a close structural analogue of galaxolide with similar physical and chemical properties, galaxolide is considered a suitable read-across analogue for determination of the hazard characteristics of musk 89.

## Introduction and use

### Australia

No specific Australian import or manufacturing information has been identified for galaxolide or musk 89. No specific Australian use has been identified for musk 89.

Galaxolide has reported domestic use in cleaning products, and in marine and automotive aftermarket products such as coatings (NICNAS, 2019). Based on information in the public domain, galaxolide is readily available for use in Australia as a fragrance ingredient (Australian Botanical Products, 2020).

### International

Available information indicates that galaxolide is used at high volumes as a fragrance ingredient in a range of products worldwide. Musk 89 does not appear to be in widespread industrial use.

Galaxolide is used in the EU in the range of 1000–10 000 tonnes per year (t/year) (REACH, 2011). In the United States of America (USA) galaxolide is listed as a high production volume

chemical (US EPA, 2020b), with an annual use volume of 454–4536 tonnes (US EPA, 2016). In the Nordic countries, the average annual use volume over a five-year period from 2014–2018 was 22 tonnes (SPIN, 2018).

Galaxolide is listed on the International Fragrance Association (IFRA) transparency list, which identifies chemicals used as fragrances by member companies (IFRA, 2020). Galaxolide is used as a fragrance ingredient in consumer products as both a base-note scent and a fixative chemical to prolong the exposure of more highly volatile fragrances (Cella and Cella, 1992; Correia, et al., 2013). Galaxolide has been measured in perfumes at concentrations up to 15 milligrams per gram (mg/g), equating to 1.5% (w/w) (Nakata, et al., 2015).

A USA-based study found and quantified galaxolide in 43 of 60 household and personal care products surveyed, with the highest concentrations found in perfumes and fragrances (up to 5.0 mg/g), body lotions and moisturisers (up to 3.7 mg/g) and deodorants (up to 2.3 mg/g) (Reiner and Kannan, 2006). Other products that contain galaxolide include body wash, shaving cream, shampoo, conditioner, hair gel, soap, cosmetics, air fresheners, scented candles, fabric softener, detergent, polishes and surface cleaners (Nakata, et al., 2015; US EPA, 2020a).

Galaxolide has non-industrial use as a fixative in pesticides to lengthen their exposure time (Besette, 2001; US EPA, 2020a). The use of this chemical as a pesticide additive is beyond the scope of this assessment.

No specific international use, import or manufacturing information has been identified for musk 89. It is listed on the European Commission's cosmetic ingredients database (CosIng) as a perfuming agent (European Commission, 2020). However, it is not registered under the EU REACH legislation and is 'inactive' on the US Environmental Protection Agency (US EPA) chemical substance inventory (ECHA, 2020; US EPA, 2020c). The chemical is not listed on the IFRA transparency list or as an ingredient in products on the chemical and product categories (CPCat) database (IFRA, 2020; US EPA, 2015). Musk 89 does not appear to have widespread industrial use internationally.

## Existing Australian regulatory controls

### Environment

The use of galaxolide and musk 89 is not subject to any specific national environmental regulations.

## International regulatory status

### United Nations

Galaxolide and musk 89 are not currently identified as persistent organic pollutants (POPs) (UNEP, 2001), ozone depleting substances (UNEP, 1987), or hazardous substances for the purpose of international trade (UNEP & FAO, 1998).

## United States of America

Galaxolide was assessed in 2014 under the US EPA's work plan. The assessment concluded that the chemical posed no unacceptable risk to the environment at current environmental exposure concentrations (US EPA, 2014).

## Environmental exposure

Galaxolide is expected to be found in household and commercial products available for use in Australia. It is used as a fragrance ingredient internationally and formulated products on the Australian market are assumed not to differ significantly from those available internationally. International studies have measured galaxolide in perfumes and fragrances, personal care products (shampoo, conditioner, soap, deodorant, body lotion, moisturiser, hair gel), and household cleaners (laundry detergent, fabric softener, surface cleaner) at concentrations up to 1.5% (w/w) (Nakata, et al., 2015; Reiner and Kannan, 2006). Chemicals used in cosmetics, personal care and cleaning products are typically released to wastewater as a normal part of their use in household and industrial applications.

Depending on degradation and partitioning processes of chemicals in sewage treatment plants (STPs), some fraction of the quantity of chemicals in wastewater entering STPs can be emitted to the air compartment, to rivers or oceans in treated effluent, or to soil by application of biosolids to agricultural land (Struijs, 1996). The emissions of galaxolide and musk 89 to environmental surface waters, sediment, and soil are considered as part of this evaluation.

## Environmental fate

### Partitioning

Galaxolide partitions to water, sediment and soil when released to the environment.

Galaxolide is a neutral organic chemical that is expected to be slightly soluble in water and moderately volatile. The Henry's Law constant of this chemical ( $36.9 \text{ Pa} \cdot \text{m}^3/\text{mol}$ ) indicates it will be moderately volatile from water and moist soil. Galaxolide is a lipophilic chemical with a measured  $\log K_{OW}$  of 5.9 and a calculated soil adsorption coefficient ( $\log K_{OC} = 4.29 \text{ L/kg}$ ), indicating that it will be immobile in soil and will preferentially adsorb to phases in the environment with high organic carbon content (including sediment and soil) (REACH, 2011; US EPA, 2017a).

Galaxolide may be emitted to the soil compartment through application of biosolids from STP processes. Calculations with a standard multimedia partitioning (fugacity) model with sole release to the soil compartment (Level III approach) predict that galaxolide will predominately be found in soil (99.9%) (US EPA, 2017a).

Galaxolide is expected to be released to the water compartment in STP effluent as a result of its use. Fugacity calculations (Level III approach) assuming sole release to the aquatic environment predict that galaxolide will primarily be found in the water (40.6%) and sediment (58.8%) compartments (US EPA, 2017a). A die-away study measured the volatilisation of galaxolide from river water and found that 17% of the initial loading volatilised over 28 days, giving a volatilisation half-life on the order of months (ECB, 2008).

Musk 89 is a neutral organic chemical that is expected to be slightly soluble in water and moderately volatile. The Henry's Law constant of this chemical (80.2 Pa·m<sup>3</sup>/mol, calc.) indicates the chemical will be moderately volatile from water and moist soil. Musk 89 is a lipophilic chemical with a calculated log K<sub>ow</sub> of 6.3 and a calculated log K<sub>oc</sub> of 4.3, which indicates it will be immobile in soil and will preferentially adsorb to phases in the environment with high organic carbon content (including sediment and soil) (US EPA, 2017a). Fugacity calculations predict that the partitioning behaviour of musk 89 will not significantly differ from galaxolide.

## Degradation

Galaxolide is persistent in some soil types. It undergoes primary degradation in water, soil and sediment to persistent degradants.

Calculations from standard QSARs predict that galaxolide will degrade rapidly in air following reaction with hydroxyl radicals with a half-life of 3.33 hours (US EPA, 2017a). Galaxolide has no hydrolysable functional groups and is not expected to degrade by hydrolysis in the water compartment.

Galaxolide is not readily biodegradable based on a study conducted according to OECD test guideline (TG) 301B (REACH, 2011). The chemical did not mineralise under screening test timeframes (0% CO<sub>2</sub> release in 28 days). A second test was conducted under the same test guidelines, using pre-adapted inoculum that had been exposed to galaxolide for 8 weeks prior. No evidence of biodegradation was observed (REACH, 2011).

Galaxolide undergoes primary degradation in water, sediment, biosolids and soil, with no evidence of ultimate degradation under environmentally relevant conditions.

The primary degradation half-life of galaxolide in river water inoculated with activated sludge was measured in a study to be 33–100 hours (ECB, 2008). After 28 days only 8% of the parent remained and the balance was accounted for by volatilisation (16%) and polar metabolites (62%). One commonly detected metabolite in the environment is galaxolidone (CAS RN 507442-49-1) (Bester, 2004; Cunha, et al., 2015; Kupper, et al., 2004; Reiner and Kannan, 2011; Vallecillos, et al., 2014). Experiments have shown that galaxolidone is not persistent itself, but it does appear to degrade more slowly than galaxolide (Poulsen and Bester, 2010). Other studies have identified methoxylated and hydroxylated metabolites of galaxolide (ECB, 2008; Martin, et al., 2007).

Galaxolide has an expected primary degradation half-life in sediment of 79 days. Sediment samples were spiked with radiolabelled galaxolide, sealed and incubated for a year. After one year, 4% of the parent galaxolide remained in the sediment, giving a primary degradation half-life of 79 days. After completion of the study there was an 80% recovery of original radioactivity from the sediment by solvent extraction, which is expected to comprise mostly polar oxidised metabolites (ECB, 2008).

Galaxolide is expected to degrade in soils with a primary degradation half-life of 95–239 days, depending on the soil type (ECB, 2008). Samples of oak forest soil, agricultural soil and STP sludge-amended agricultural soil were spiked with radiolabelled galaxolide, sealed in flasks and incubated for a year. The degradation half-lives of galaxolide were determined to be 95, 239, and 105 days for oak forest soil, agricultural soil, and sludge-amended soil, respectively. In the sludge-amended soil the total recovery of radiolabelled material was 73%, comprising galaxolide (9%) and polar degradants. An additional 20% of radiolabelled material was recovered after treatment by a process to detach covalently bound degradants

from organic components in the soil. Therefore, galaxolide may degrade in soil with a half-life exceeding the domestic categorisation threshold for persistence (EPHC, 2009).

Studies with pure fungal cultures show that some mineralisation may occur under idealised and not environmentally relevant conditions. A study was conducted to isolate highly active fungal strains from 64 soil samples, and resulting pure fungal cultures were tested for their capacity to degrade radiolabelled galaxolide. In cultures of *Phanerochaete chrysosporium* (a white rot fungus) galaxolide disappeared within 3 days, whereas in *Cladosporium cladosporioides* (one of the most common fungi in outdoor air world-wide) 95% of galaxolide disappeared within 4 weeks (ECB, 2008). The detected metabolites were more polar than the parent and comprised galaxolidone (19%) and other uncharacterised degradants (75%). Evidence of partial mineralisation to CO<sub>2</sub> was observed after an extended period, with 82% of total radioactivity recovered from organic and aqueous extractions, and 18% recovered as CO<sub>2</sub> after 200 days. The mineralisation half-life under these idealised conditions would, therefore, still exceed domestic categorisation thresholds (EPHC, 2009).

No studies on the degradation of musk 89 have been identified but the chemical is expected to be persistent based on standard QSARs (US EPA, 2017a), with predicted primary and ultimate half-lives of 7.6 months and more than 10 years, respectively (LMC, 2015).

## Bioaccumulation

Galaxolide and musk 89 have a moderate potential to bioaccumulate in aquatic life, and an uncertain potential to bioaccumulate in benthic organisms. Metabolites of galaxolide are more polar than the parent chemical and are not expected to bioaccumulate.

Experimentally determined bioconcentration factors (BCFs) are below the domestic categorisation threshold for bioaccumulation (BCF  $\geq$  2000 L/Kg) (EPHC, 2009). Two bioconcentration studies have been conducted in accordance with OECD TG 305E; a BCF of 1584 litres per kilogram (L/Kg) wet weight (wwt) was reported for *Lepomis macrochirus* (bluegill sunfish) (REACH, 2011), and a BCF of 1660 L/kg wwt was reported for *Danio rerio* (Balk, et al., 2001). The BCF for the latter study was calculated from the lipid-only BCF, normalised to 5% lipid weight. Non-standard studies in the scientific literature report BCFs in the range of 201–1561 L/kg wwt (Blum, et al., 2018; Fromme, et al., 2001; Schreurs, et al., 2004), and bioaccumulation factors (BAFs) in the range 18–371 L/Kg wwt (Reiner and Kannan, 2011).

There is some evidence to suggest that galaxolide has a higher bioaccumulation potential in some sediment-dwelling organisms. One study exposed *Lumbriculus variegatus* (blackworm) to galaxolide in a water-only test vessel, deriving a BCF value of 2692 L/Kg wwt (Artola-Garicano, et al., 2003). Tests were also conducted with *Chironomus riparius* larvae (midge), deriving a BCF of 85 L/kg. Since these tests were not conducted in mixed sediment-water systems it is not possible to conclude if these bioconcentration values are environmentally relevant.

There are limited data available to evaluate the bioaccumulation potential of the metabolites of galaxolide. However, metabolites are each expected to be more polar than galaxolide, and may be cleared from aquatic organisms quite rapidly (ECB, 2008; Fernandes, et al., 2013; Martin, et al., 2007). During the principal OECD TG 305E bioconcentration study for galaxolide with *L. macrochirus*, it was estimated that galaxolide was metabolised and the metabolite(s) excreted with a turnover rate of 38–50% per day (ECB, 2008).

Musk 89 is not expected to be bioaccumulative based on calculations with standard QSPRs that predict a BCF of 1156 L/kg, which is below the domestic categorisation threshold.

## Environmental transport

Galaxolide may undergo long-range transport. Galaxolide has been identified in remote pristine environmental areas in biota, air, surface water and snow samples.

Despite a short predicted atmospheric lifetime of 3.3–5.3 hours (Aschmann, et al., 2001), environmental monitoring has identified galaxolide in pristine environmental areas that do not receive direct anthropogenic outputs. Galaxolide was found in fish caught at remote alpine lakes in Switzerland in concentrations of 0.042–0.078 milligrams per kilogram (mg/kg) lipid weight (Schmid, et al., 2007). Atmospheric concentrations of galaxolide were measured in the North Sea and Arctic regions in concentrations ranging from 5–49 picograms per cubic metre (pg/m<sup>3</sup>) and 1–9 pg/m<sup>3</sup>, respectively. Surface water concentrations in the North Sea ranged from 12–2030 picograms per litre (pg/L), with quantities in the ocean decreasing with distance from the mainland in a north/northeast direction (none detected in samples from the northeast Atlantic or Arctic) (Xie, et al., 2007). Galaxolide has also been measured in snow samples collected from a remote glacier in the Italian Alps, in concentrations of 2.13–5.07 ng/L (Villa, et al., 2014).

## Predicted environmental concentration (PEC)

The concentration of galaxolide in Australian river water is estimated to be 2.0 micrograms per litre (µg/L), based on domestic monitoring data, with a worst-case concentration of 4.6 µg/L. Reasonable worst-case concentrations of galaxolide in domestic sediments and soil are predicted to be 0.38 mg/kg dry weight (dw) and 0.11 mg/kg dw respectively. These values were determined by considering available domestic and international monitoring data for galaxolide in wastewater effluents, biosolids, sediments, surface waters and biota. No environmental monitoring data for musk 89 were identified, and a PEC has not been determined.

The highest galaxolide concentrations in the Australian environment were measured at sewage outfall sites in Darwin Harbour in 2010–2011. The maximum concentration measured was 2.0 µg/L, detected in the outfall from East Point STP in October 2010 (late dry season), an advanced primary treatment plant that treats urban sewage from Darwin city. Treated effluent is released 150 m offshore into Darwin Harbour, and samples were collected near the discharge pipe at low tide to minimise dilution. Concentrations of 0.1–0.5 µg/L were measured at other sites near Darwin at different times of the year (French, et al., 2015). The value of 2.0 µg/L in surface water is taken as the estimated galaxolide surface water concentration in Australia for the purposes of risk characterisation.

A recent domestic study tracked galaxolide through an advanced water treatment plant in Sydney, NSW (Wang and Khan, 2014). The study reported an average influent concentration of 2.55 µg/L that reduced to 1.19 µg/L after secondary treatment (moving bed biofilm reactor and membrane bioreactor), and 0.021 µg/L after tertiary treatment (reverse osmosis and disinfection, 99% total removal efficiency).

Galaxolide has also been detected in STP influent and effluent internationally. The removal efficiency following secondary and tertiary treatment is consistent with measured values in Australia. Influent concentrations up to 45.1 µg/L have been measured internationally (Vallecillos, et al., 2014), but values are more frequently in the range of 0.5–10 µg/L (Bester, 2004; Clara, et al., 2011; Ramirez, et al., 2011; Simonich, et al., 2002; Smyth, et al., 2007). Effluent concentrations largely depend on the level of treatment available, with concentrations of ca 0.2–4.6 µg/L measured following secondary treatment (Bester, 2004; Buerge, et al., 2003; Clara, et al., 2011; Ramirez, et al., 2011; Simonich, et al., 2002; Smyth, et al., 2007; Vallecillos, et al., 2014; Wong, et al., 2019) and concentrations of 0.0015–0.042

µg/L measured following tertiary treatment (Ramírez, et al., 2011; Vallecillos, et al., 2014). The value of 4.6 µg/L in effluent after secondary treatment is taken as a worst-case surface water concentration in Australia.

International studies have quantified galaxolide in surface waters (rivers and lakes) in concentrations up to 3.15 µg/L (Blum, et al., 2018; Buerge, et al., 2003; Clara, et al., 2011; Fromme, et al., 2001; Oros, et al., 2003; Ramírez, et al., 2011; Wong, et al., 2019). An extensive analysis of surface waters in Germany reported a 90<sup>th</sup> percentile galaxolide concentration of 0.46 µg/L (OSPAR Commission, 2004). Concentrations tend to be higher in urban area waterways and those impacted by sewage outfall (Fromme, et al., 2001; Wong, et al., 2019).

A study of sediment contamination in Berlin (Germany) rivers reported a 90<sup>th</sup> percentile galaxolide concentration of 0.38 mg/kg dw from a river moderately contaminated with biosolids, and a maximum concentration of 2.20 mg/kg from a heavily contaminated river (Fromme, et al., 2001). Galaxolide has been measured in concentrations up to 0.27 mg/kg dw in sediment samples from Europe (Sweden and Austria) and China (Pearl River estuary) (Blum, et al., 2018; Clara, et al., 2011; Huang, et al., 2016). A value of 0.38 mg/kg dw has been used as the concentration of galaxolide in sediments in Australia for the purposes of risk characterisation and 2.20 mg/kg dw is considered a worst-case PEC for the purposes of this evaluation.

Measured concentrations of galaxolide in biosolids range of 0.25–177 mg/kg dw internationally (Bester, 2004; Clara, et al., 2011; DiFrancesco, et al., 2004; Kupper, et al., 2004; Langdon, et al., 2010; Mogensen, et al., 2004). Mean and median values of 14.0 and 10.7 mg/kg dw, respectively, were determined in a literature review of biosolids contaminants (Langdon, et al., 2010).

The calculated galaxolide concentration in soil amended with biosolids is 0.11 mg/kg dw based on mean measured international biosolid concentrations (14 mg/kg dw), typical biosolids application rates and a soil bulk density of 1300 kilograms per cubic metre (kg/m<sup>3</sup>) (EPHC, 2009; Langdon, et al., 2010). A maximum concentration of 5.5 mg/kg dw was calculated using the maximum measured concentration of galaxolide in biosolids and maximum biosolids application rates.

Galaxolide has been found in aquatic biota internationally. Galaxolide has been detected in perch (0.0067–0.13 mg/kg wwt, 0.69–27 mg/kg lipid weight) (Blum, et al., 2018; Törneman and SWECO Environment, 2008), eel (up to 4.8 mg/kg wwt, 18.48 mg/kg lipid weight) (Fromme, et al., 2001), mullet and flounder (up to 0.012 mg/kg dw), mussels (up to 0.035 mg/kg dw) and clams (0.033 mg/kg dw) (Cunha, et al., 2015; Törneman and SWECO Environment, 2008). A study by Fromme and co-workers correlated the concentration of galaxolide in eels with the extent of river contamination from STP effluent. Areas with low, medium and high contamination had average concentrations of 0.052, 0.12 and 1.51 mg/kg wwt, respectively (Fromme, et al., 2001).

Environmental concentrations of musk 89 have not been estimated using default introduction volumes. Based on available information, musk 89 does not have widespread industrial use. If musk 89 is used as a fragrance ingredient domestically it is only expected to be used in small volumes resulting in limited environmental emissions.

# Environmental effects

## Effects on aquatic life

Galaxolide and musk 89 are expected to cause toxic effects at low concentrations in aquatic organisms across multiple trophic levels.

No experimental data for the ecotoxicity of musk 89 were identified. The ecotoxicity data presented below for galaxolide is expected to be suitable for read-across to musk 89 given the structural similarity of the two chemicals and expected baseline narcosis mode of toxic action for both chemicals (LMC, 2020; US EPA, 2017b).

### Acute toxicity

The following measured median lethal concentration (LC50) and median effective concentration (EC50) values for model organisms across three trophic levels exposed to galaxolide were retrieved from the scientific literature or the registration dossier for galaxolide under EU REACH legislation (REACH, 2011; Yamauchi, et al., 2008):

Taxon	Endpoint	Method
Fish	96 h LC50 = 0.95 mg/L	<i>Oryzias latipes</i> larvae (Japanese rice fish) Test substance was dissolved in 0.1% DMSO OECD TG 203 equivalent
	96 h LC50 = 1.36 mg/L	<i>Lepomis macrochirus</i> (bluegill sunfish) Flow-through conditions OECD TG 204
Invertebrate	48 h EC50 = 0.3 mg/L	<i>Daphnia magna</i> (water flea) Mobility OECD TG 202
Algae	72 h EC50 = 0.72 mg/L	<i>Pseudokirchneriella subcapitata</i> (microalgae) Biomass OECD TG 201

### Chronic toxicity

The following measured no-observed-effect concentrations (NOEC) for model organisms across three trophic levels were retrieved from the scientific and regulatory literature, or the registration dossier for galaxolide under EU REACH legislation (Balk and Ford, 1999; REACH, 2011):

Taxon	Endpoint	Method
Fish	36 d NOEC = 0.068 mg/L	<i>Pimephales promelas</i> eggs (fathead minnow) Larval survival and growth Flow-through conditions OECD TG 210
	21 d NOEC = 0.093 mg/L	<i>Lepomis macrochirus</i> (bluegill sunfish) Growth Flow-through conditions OECD TG 204 (extended to 21 d)
Invertebrate	21 d NOEC = 0.111 mg/L	<i>Daphnia magna</i> (water flea) Reproduction Semi-static conditions OECD TG 202
Algae	72 h NOEC = 0.201 mg/L	<i>Pseudokirchneriella subcapitata</i> (microalgae) Growth Static conditions OECD TG 201

### Marine chronic toxicity

The following measured effective concentration at 10% inhibition (EC10) value for a model invertebrate organism exposed to galaxolide were retrieved from the registration dossier for the chemical under EU REACH legislation (REACH, 2011):

Taxon	Endpoint	Method
Invertebrate	5–6 d EC10 = 0.0438 mg/L	<i>Acartia tonsa</i> (marine copepod) Larval development rate OECD Draft Guideline

Galaxolide saturates efflux transporters in the gills of the marine mussel *Mytilus californianus*, impeding their ability to eject xenobiotic substances from cells and raising concern that galaxolide might enhance the toxicity of other chemicals. Inhibition was observed for 24–48 hours following a two-hour exposure of galaxolide, with a median inhibitory concentration (IC50) of 0.63 mg/L (2.43 µM) (Luckenbach and Epel, 2005). Insufficient data are currently available to evaluate the potential for this effect to cause adverse outcomes in organisms exposed to galaxolide in the environment.

### Effects on sediment dwelling life

Galaxolide can cause toxic effects in sediment-dwelling organisms at low concentrations.

## Sediment-dwelling acute toxicity

The following measured LC50 value for amphipods exposed to galaxolide in sediment were retrieved from the Registration Dossier for the chemical under EU REACH legislation (REACH, 2011):

Taxon	Endpoint	Method
Amphipod	96 h LC50 = 62.5 mg/kg dw	<i>Hyalella azteca</i> (scud) OECD TG 218

## Sediment-dwelling chronic toxicity

The following NOEC values for three model organisms exposed to galaxolide in sediments were retrieved from the Registration Dossier for the chemical under EU REACH legislation (REACH, 2011):

Taxon	Endpoint	Method
Midge	28 d NOEC = 250 mg/kg dw	<i>Chironomus riparius</i> (harlequin fly) Emergence rate OECD TG 218
Worm	28 d NOEC = 16.2 mg/kg dw	<i>Lumbriculus variegatus</i> (blackworm) Reproduction OECD TG 218
Amphipod	28 d NOEC = 7.1 mg/kg dw	<i>Hyalella azteca</i> (scud) Growth OECD TG 218

## Effects on terrestrial life

Galaxolide can cause toxic effects in terrestrial organisms.

Chronic ecotoxicity values have been obtained for the springtail *Folsomia candida* and the earthworm *Eisenia fetida* exposed to galaxolide in standard soil with 10% added organic matter (sphagnum peat). Both studies were conducted according to ISO test guidelines. A 28 day NOEC of 45 mg/kg dw (survival) was obtained for *F. candida*, and for *E. fetida* a 56 day NOEC (reproduction) of 45 mg/kg dw and a 28 day NOEC (growth) of 105 mg/kg dw were obtained (REACH, 2011).

## Endocrine effects/activity

Galaxolide has some endocrine activity at the biochemical level in fish. However, no apical effects have been observed as a result of endocrine activity.

Galaxolide exposure at 0.5 mg/L induced higher levels of hepatic vitellogenin in medaka (*Oryzias latipes*), indicating some estrogenic activity (Yamauchi, et al., 2008). Galaxolide has also exhibited anti-estrogenic activity; when co-injected with estradiol in rainbow trout (*Oncorhynchus mykiss*), plasma levels of vitellogenin decreased (Simmons, et al., 2010). In

vitro studies have also measured an anti-estrogenic response against fish estrogen receptors (Schreurs, et al., 2004; Simmons, et al., 2010).

## Predicted no-effect concentration (PNEC)

A freshwater PNEC for galaxolide of 6.8 µg/L was derived from the measured fish chronic ecotoxicity endpoint (36 d NOEC = 0.068 mg/L), using an assessment factor of 10. This assessment factor was selected as reliable chronic ecotoxicity data are available over three trophic levels (EPHC, 2009).

A soil PNEC of 0.31 mg/kg dw was derived from the measured springtail chronic ecotoxicity endpoint (28 d NOEC = 45 mg/kg dw). Following the approach used in the EU Risk Assessment Report, the NOEC for *F. candida* was first normalised to a soil organic content of 3.4% (ECB, 2008). An assessment factor of 50 was then applied, selected because reliable chronic ecotoxicity data are available over two taxa (EPHC, 2009).

A sediment PNEC of 1.58 mg/kg dw was derived from the measured midge chronic ecotoxicity endpoint (28 d NOEC = 7.1 mg/kg dw). Following the approach above, the NOEC for *H. azteca* was first normalised to a sediment organic content of 4% (EPHC, 2009). An assessment factor of 10 was then applied, selected because reliable chronic ecotoxicity data are available over three taxa.

## Categorisation of environmental hazard

The categorisation of the environmental hazards of the assessed chemicals according to domestic environmental hazard thresholds is presented below (EPHC, 2009):

### Persistence

Persistent (P). Based on results from standard biodegradability tests that show no evidence of ultimate degradation and evidence that galaxolide degrades to persistent metabolites, galaxolide and musk 89 are categorised as Persistent in sediment and water. Based on results from degradation studies that report a primary half-life that exceeds 180 days in some soil types, galaxolide and musk 89 are categorised as Persistent in soil.

### Bioaccumulation

Not Bioaccumulative (Not B). Based on low measured bioconcentration factors (BCF) in fish, evidence of biotransformation, and low potential for metabolites to bioaccumulate, galaxolide and musk 89 are categorised as Not Bioaccumulative.

### Toxicity

Toxic (T). Based on available acute ecotoxicity values below 1 mg/L and evidence of chronic toxicity (ecotoxicity values below 0.1 mg/L), galaxolide and musk 89 are categorised as Toxic.

## Environmental risk characterisation

Based on the PEC and PNEC values determined above, the following Risk Quotients (RQ = PEC ÷ PNEC) have been calculated for release of galaxolide into rivers, soil and sediment:

Compartment	PEC	PNEC	RQ
River	2.0 µg/L	6.8 µg/L	0.29
Soil	0.11 mg/kg dw	0.31 mg/kg dw	0.35
Sediment	0.38 mg/kg dw	1.58 mg/kg dw	0.24

For rivers, an RQ less than 1 indicates that galaxolide is not expected to pose a high risk to the environment based on estimated emissions, as environmental concentrations are below levels likely to cause harmful effects. The RQ considering the worst-case PEC of 4.6 µg/L is 0.68, indicating that galaxolide is not expected to pose a high risk to the environment even at worst-case concentrations in the freshwater aquatic compartment.

For soil, an RQ less than 1 indicates that galaxolide is not expected to pose a high risk to the environment based on estimated emissions, as environmental concentrations are below levels likely to cause harmful effects. The RQ considering the worst-case PEC of 5.5 mg/kg dw is 17.7. This value is calculated using the highest recorded concentration of galaxolide in biosolids in an international study, and concentrations of these magnitudes are unlikely to occur in Australia. However, this worst-case scenario indicates that galaxolide could pose a high risk to the environment if severely contaminated biosolids are applied to agricultural soils in Australia.

For sediment, an RQ less than 1 indicates that galaxolide is not expected to pose a high risk to the environment based on estimated emissions, as environmental concentrations are below levels likely to cause harmful effects. The RQ considering the worst-case PEC of 2.20 mg/kg dw is 1.4. This PEC is the highest recorded concentration of galaxolide in sediment in an international study, from a heavily contaminated urban river in Berlin, Germany. Concentrations of these magnitudes are unlikely to occur in Australia. However, an RQ above 1 indicates that galaxolide could pose a high risk to the environment at worst-case concentrations in sediment.

Marine aquatic organisms may be more sensitive to galaxolide than freshwater organisms of the same trophic level, though there are fewer studies available. However, emissions of galaxolide to the marine environment are unlikely to be a concern where discharge occurs into well-mixed marine waters.

Musk 89 is expected to have similar hazard characteristics, but much lower environmental exposures, to galaxolide. Given the determination that current releases of galaxolide are unlikely to be a concern, the much lower release volumes anticipated for musk 89 mean that this chemical is also unlikely to be a concern. It is therefore not expected to pose a high risk to the environment in water, sediment or soil.

## Uncertainty

This evaluation was conducted based on a set of information that may be incomplete or limited in scope. Some relatively common data limitations can be addressed through use of conservative assumptions (OECD, 2019) or quantitative adjustments such as assessment factors (OECD, 1995). Others must be addressed qualitatively, or on a case-by-case basis (OECD, 2019).

The most consequential areas of uncertainty for this evaluation are:

- The identity, environmental fate and effects of recalcitrant degradants of galaxolide are unknown. If information becomes available in the future to indicate that these degradants may have persistent, bioaccumulative and toxic (PBT) characteristics, the PBT categorisation of galaxolide may also change.
- There are no domestic monitoring data for galaxolide in the sediment or soil compartments. Measured international concentrations in these compartments indicate that galaxolide may be present at concentrations exceeding the level of concern in heavily contaminated soils and sediments. The risk profile of galaxolide may change should monitoring data become available to indicate that galaxolide is present in Australian soils or sediments at levels above the levels of concern.
- This evaluation focuses on the discrete chemical known as galaxolide (CAS RN 1222-05-5), but in practice galaxolide is used as a technical mixture containing minor structural isomers in concentrations up to 30%. Information was not readily available on whether the test substance used in standard tests (physical and chemical properties, biodegradation, bioaccumulation and toxicity) was pure galaxolide, or the technical mixture containing significant amounts of the structural isomers of galaxolide. It is also uncertain how each of these materials would impact the results of each test, though it is assumed that the properties of the structural isomers are closely modelled by galaxolide. The risk profile of galaxolide may change if information becomes available to indicate that the structural isomers of galaxolide present as impurities in the technical mixture have more hazardous properties than galaxolide.

## References

Artola-Garicano E, Sinnige TL, van Holsteijn I, Vaes WHJ and Hermens JLM (2003). Bioconcentration and acute toxicity of polycyclic musks in two benthic organisms (*Chironomus riparius* and *Lumbriculus variegatus*). *Environmental Toxicology and Chemistry*, **22**(5), pp 1086-1092.

Aschmann SM, Arey J, Atkinson R and Simonich SL (2001). Atmospheric Lifetimes and Fates of Selected Fragrance Materials and Volatile Model Compounds. *Environmental Science & Technology*, **35**(18), pp 3595-3600.

Australian Botanical Products (2020). *Aromatic chemicals list*. Accessed October 2020 at <https://www.abp.com.au/files/Aroma%20Chem%20List%20Brochure.pdf>.

Balk F and Ford RA (1999). Environmental risk assessment for the polycyclic musks, AHTN and HHCB: II. Effect assessment and risk characterisation. *Toxicology Letters*, **111**(1), pp 81-94.

Balk F, Blok H and Salvito D (2001). Environmental Risks of Musk Fragrance Ingredients. In: ed. *Pharmaceuticals and Care Products in the Environment*. American Chemical Society, pp 168-190.

Bessette SM (2001). *Pesticidal compositions containing plant essential oils against termites (WO2001000020)*. Accessed November 2020 at <https://patents.google.com/patent/WO2001000020A1/en?q=WO2001000020>.

Bester K (2004). Retention characteristics and balance assessment for two polycyclic musk fragrances (HHCB and AHTN) in a typical German sewage treatment plant. *Chemosphere*, **57**(8), pp 863-870.

- Blum KM, Andersson PL, Ahrens L, Wiberg K and Haglund P (2018). Persistence, mobility and bioavailability of emerging organic contaminants discharged from sewage treatment plants. *Science of The Total Environment*, **612**, pp 1532-1542.
- Buerge IJ, Buser H-R, Müller MD and Poiger T (2003). Behavior of the Polycyclic Musks HHCb and AHTN in Lakes, Two Potential Anthropogenic Markers for Domestic Wastewater in Surface Waters. *Environmental Science & Technology*, **37**(24), pp 5636-5644.
- Cella JA and Cella, JA, III (1992). *Method for enhancing fragrance applications (US5120709A)*. Accessed September October 2020 at <https://patents.google.com/patent/US5120709A/en?q=US+5120709>.
- Clara M, Gans O, Windhofer G, Krenn U, Hartl W, Braun K, Scharf S and Scheffknecht C (2011). Occurrence of polycyclic musks in wastewater and receiving water bodies and fate during wastewater treatment. *Chemosphere*, **82**(8), pp 1116-1123.
- Correia P, Cruz A, Santos L and Alves A (2013). Human dermal exposure to galaxolide from personal care products. *International Journal of Cosmetic Science*, **35**(3), pp 299-309.
- Cunha SC, Fernandes JO, Vallecillos L, Cano-Sancho G, Domingo JL, Pocurull E, Borrull F, Maulvault AL, Ferrari F, Fernandez-Tejedor M, Van den Heuvel F and Kotterman M (2015). Co-occurrence of musk fragrances and UV-filters in seafood and macroalgae collected in European hotspots. *Environmental Research*, **143**, pp 65-71.
- DiFrancesco AM, Chiu PC, Standley LJ, Allen HE and Salvito DT (2004). Dissipation of Fragrance Materials in Sludge-Amended Soils. *Environmental Science & Technology*, **38**(1), pp 194-201.
- ECB (2008). *European Union Risk Assessment Report: 1,3,4,6,7,8-hexahydro-4,6,6,7,8,8-hexamethylcyclopenta-γ-2-benzopyran (CAS No: 1222-05-5)*. European Chemicals Bureau, The Netherlands. Accessed September 2020 at <https://echa.europa.eu/documents/10162/947def3b-bbbf-473b-bc19-3bda7a8da910>.
- ECHA (2020). *Registered Substances List*. European Chemicals Agency, Helsinki, Finland. Accessed December 2020 at <https://echa.europa.eu/information-on-chemicals/registered-substances>.
- EPHC (2009). *Environmental Risk Assessment Guidance Manual for Industrial Chemicals*. Environment Protection and Heritage Council (EPHC), Canberra, Australia. Accessed September 2020 at <http://www.nepc.gov.au/resource/chemical-risk-assessment-guidance-manuals>.
- European Commission (2020). *Cosmetic Ingredient Database (CosIng) Ingredient: hexahydro-4,6,6,9,9-pentamethyl-naphthopyran (CAS RN 1922-67-4)*. European Commission, Brussels, Belgium. Downloaded Available at [https://ec.europa.eu/growth/tools-databases/cosing/index.cfm?fuseaction=search.details\\_v2&id=40164](https://ec.europa.eu/growth/tools-databases/cosing/index.cfm?fuseaction=search.details_v2&id=40164).
- Fernandes D, Dimastrogiovanni G, Blázquez M and Porte C (2013). Metabolism of the polycyclic musk galaxolide and its interference with endogenous and xenobiotic metabolizing enzymes in the European sea bass (*Dicentrarchus labrax*). *Environmental Pollution*, **174**, pp 214-221.
- French VA, Codi King S, Kumar A, Northcott G, McGuinness K and Parry D (2015). Characterisation of microcontaminants in Darwin Harbour, a tropical estuary of northern

Australia undergoing rapid development. *Science of The Total Environment*, **536**, pp 639-647.

Fromme H, Otto T and Pilz K (2001). Polycyclic musk fragrances in different environmental compartments in Berlin (Germany). *Water Research*, **35**(1), pp 121-128.

Heeringa LG and Beets MGJ (1963). *Tricyclic isochromans and processes for making same*. International Flavors & Fragrances Inc., New York, USA. Accessed September 2020 at <https://patents.google.com/patent/US3360530A/en?q=US3360530>.

HERA (2004). *Human and Environmental Risk Assessment on ingredients of household cleaning products: HHCB (1,3,4,6,7,8-hexahydro-4,6,6,7,8,8-hexamethylcyclopenta-γ-2-benzopyran and related isomers)*. The HERA (Human and Environmental Risk Assessment) Project, Belgium, Brussels. Accessed October 2020 at <https://www.heraproject.com/files/29-HH-04-pcm%20HHCB%20HERA%20Human%20Health%20DISCL%20ed2.pdf>.

Huang W, Xie Z, Yan W, Mi W and Xu W (2016). Occurrence and distribution of synthetic musks and organic UV filters from riverine and coastal sediments in the Pearl River estuary of China. *Marine Pollution Bulletin*, **111**(1), pp 153-159.

IFRA (2020). *International Fragrance Association (IFRA) Transparency List*. International Fragrance Association Geneva, Switzerland. Downloaded September 2020. Available at <https://ifrafragrance.org/initiatives/transparency/ifra-transparency-list>.

Kupper T, Berset JD, Etter-Holzer R, Furrer R and Tarradellas J (2004). Concentrations and specific loads of polycyclic musks in sewage sludge originating from a monitoring network in Switzerland. *Chemosphere*, **54**(8), pp 1111-1120.

Langdon K, Warne M and Kookana R (2010). Aquatic hazard assessment for pharmaceuticals, personal care products, and endocrine-disrupting compounds from biosolids-amended land. *Integrated Environmental Assessment and Management*, **6**(4), pp 663-76.

LMC (2015). *OASIS Catalogic*, v 5.11.17. Laboratory of Mathematical Chemistry (LMC), University "Prof. Dr. Assen Zlatarov", Burgas, Bulgaria. Available at <http://oasis-lmc.org/>.

LMC (2020). *The OECD QSAR Toolbox*, v 4.4.1. Laboratory of Mathematical Chemistry (LMC), University "Prof. Dr. Assen Zlatarov", Burgas, Bulgaria. Available at <https://qsartoolbox.org/>.

Luckenbach T and Epel D (2005). Nitromusk and polycyclic musk compounds as long-term inhibitors of cellular xenobiotic defense systems mediated by multidrug transporters. *Environmental health perspectives*, **113**(1), pp 17-24.

Martin C, Moeder M, Daniel X, Krauss G and Schlosser D (2007). Biotransformation of the Polycyclic Musks HHCB and AHTN and Metabolite Formation by Fungi Occurring in Freshwater Environments. *Environmental Science & Technology*, **41**(15), pp 5395-5402.

Mogensen BB, Pritzl G, Rastogi S, Glesne O, Hedlund B, Hirvi J-P, Lundgren A and Sigurdsson A (2004). *Musk Compounds in the Nordic Environment*. TemaNord, Copenhagen, Denmark.

Nakata H, Hinosa M and Yanagimoto H (2015). Macrocyclic-, polycyclic-, and nitro musks in cosmetics, household commodities and indoor dusts collected from Japan: Implications for their human exposure. *Ecotoxicology and Environmental Safety*, **111**, pp 248-255.

NICNAS (2016). *Tonalide and related polycyclic musks: Environment tier II assessment*. National Industrial Chemicals Notification and Assessment Scheme, Sydney, Australia. Accessed October 2020 at <https://www.industrialchemicals.gov.au/sites/default/files/Tonalide%20and%20related%20polycyclic%20musks%20Environment%20tier%20II%20assessment.pdf>.

NICNAS (2017). *Celestolide and related polycyclic musks: Environment tier II assessment*. National Industrial Chemicals Notification and Assessment Scheme Sydney, Australia. Accessed October 2020 at <https://www.industrialchemicals.gov.au/sites/default/files/Celestolide%20and%20related%20polycyclic%20musks%20Environment%20tier%20II%20assessment.pdf>.

NICNAS (2019). *Cyclopenta[g]-2-benzopyran, 1,3,4,6,7,8-hexahydro-4,6,5,7,8,8-hexamethyl-: Human health tier II assessment*. National Industrial Chemicals Notification and Assessment Scheme, Sydney, Australia. Accessed February 2021 at <https://www.industrialchemicals.gov.au/sites/default/files/Cyclopenta%5Bg%5D-2-benzopyran%2C%201%2C3%2C4%2C6%2C7%2C8-hexahydro-4%2C6%2C6%2C7%2C8%2C8-hexamethyl-Human%20health%20tier%20II%20assessment.pdf>.

OECD (1995). *Guidance document for aquatic effects assessment*. Organisation for Economic Co-operation and Development, Paris.

OECD (2019). *Guiding Principles and Key Elements for Establishing a Weight of Evidence for Chemical Assessment, Series on Testing and Assessment No. 311, Environment, Health and Safety Division, Environment Directorate* The Organisation for Economic Co-operation and Development Accessed April 2021 at <https://www.oecd.org/chemicalsafety/risk-assessment/guiding-principles-and-key-elements-for-establishing-a-weight-of-evidence-for-chemical-assessment.pdf>.

Oros DR, Jarman WM, Lowe T, David N, Lowe S and Davis JA (2003). Surveillance for previously unmonitored organic contaminants in the San Francisco Estuary. *Marine Pollution Bulletin*, **46**(9), pp 1102-1110.

OSPAR Commission (2004). Hazardous Substances Series, *Musk xylene and other musks*. Convention for the Protection of the Marine Environment of the North-East Atlantic (OSPAR Convention) Commission, London, United Kingdom. Accessed October 2020 at <https://www.ospar.org/documents?v=6978>.

Poulsen TG and Bester K (2010). Organic Micropollutant Degradation in Sewage Sludge during Composting under Thermophilic Conditions. *Environmental Science & Technology*, **44**(13), pp 5086-5091.

Ramírez N, Marcé RM and Borrull F (2011). Development of a stir bar sorptive extraction and thermal desorption–gas chromatography–mass spectrometry method for determining synthetic musks in water samples. *Journal of Chromatography A*, **1218**(1), pp 156-161.

REACH (2011). *Registration, Evaluation, Authorisation and Restriction of Chemicals (REACH) Dossier for 1,3,4,6,7,8-hexahydro-4,6,6,7,8,8-hexamethylindeno[5,6-c]pyran*.

European Chemicals Agency. Accessed September 2020 at <https://echa.europa.eu/registration-dossier/-/registered-dossier/14504>.

Reiner JL and Kannan K (2006). A survey of polycyclic musks in selected household commodities from the United States. *Chemosphere*, **62**(6), pp 867-873.

Reiner JL and Kannan K (2011). Polycyclic Musks in Water, Sediment, and Fishes from the Upper Hudson River, New York, USA. *Water, Air, & Soil Pollution*, **214**(1), pp 335-342.

Schmid P, Kohler M, Gujer E, Zennegg M and Lanfranchi M (2007). Persistent organic pollutants, brominated flame retardants and synthetic musks in fish from remote alpine lakes in Switzerland. *Chemosphere*, **67**(9), pp S16-S21.

Schreurs RHMM, Legler J, Artola-Garicano E, Sinnige TL, Lanser PH, Seinen W and van der Burg B (2004). In Vitro and in Vivo Antiestrogenic Effects of Polycyclic Musks in Zebrafish. *Environmental Science & Technology*, **38**(4), pp 997-1002.

Simmons DBD, Marlatt VL, Trudeau VL, Sherry JP and Metcalfe CD (2010). Interaction of Galaxolide® with the human and trout estrogen receptor- $\alpha$ . *Science of The Total Environment*, **408**(24), pp 6158-6164.

Simonich SL, Federle TW, Eckhoff WS, Rottiers A, Webb S, Sabaliunas D and de Wolf W (2002). Removal of Fragrance Materials during U.S. and European Wastewater Treatment. *Environmental Science & Technology*, **36**(13), pp 2839-2847.

Smyth SA, Lishman LA, McBean EA, Kleywegt S, Yang J-J, Svoboda ML, Ormonde S, Pileggi V, Lee H-B and Seto P (2007). Polycyclic and Nitro Musks in Canadian Municipal Wastewater: Occurrence and Removal in Wastewater Treatment. *Water Quality Research Journal*, **42**(3), pp 138-152.

SPIN (2018). *Substance record: 1,3,4,6,7,8-hexahydro-4,6,6,7,8,8-hexamethylindeno[5,6-c]pyran (CAS RN 1222-05-5)*. Substances in Preparations in Nordic Countries (SPIN), København, Denmark. Downloaded August 2020. Available at <http://www.spin2000.net/spinmyphp/>.

Sprecker MA (1987). *Single phase liquid mixture of tricyclic isochroman derivative mixture and acetyl tetrahydronaphthalene derivative mixture*. International Flavors and Fragrances Inc., New York, USA. Accessed October 2020 at <https://patents.google.com/patent/US4650603A/en?q=US4650603>.

Struijs J (1996). *SimpleTreat 3.0: a model to predict the distribution and elimination of chemicals by sewage treatment plants*. National Institute of Public Health and the Environment, Bilthoven, The Netherlands.

Törneman N and SWECO Environment, AB, (2008). SWECO Environment Screening Report 2008:2, *Screening of musk substances*. Malmö, Sweden.

UNEP (1987). *The Montreal Protocol on Substances that Deplete the Ozone Layer*. United Nations Environment Programme, Ozone Secretariat, Nairobi, Kenya. Accessed October 2020 at <http://ozone.unep.org/>.

UNEP (2001). *The Stockholm Convention on Persistent Organic Pollutants*. United Nations Environment Programme, Secretariat of the Stockholm Convention, Châtelaine, Switzerland. Accessed October 2020 at <http://www.pops.int/>.

UNEP & FAO (1998). *The Rotterdam Convention on the Prior Informed Consent Procedure for Certain Hazardous Chemicals and Pesticides in International Trade*. United Nations Environment Programme and Food and Agriculture Organization of the United Nations, Châtelaine, Switzerland. Accessed October 2020 at <http://www.pic.int/>.

US EPA (2014). TSCA Work Plan Chemical Risk Assessment, *HHCB, 1,3,4,6,7,8-hexahydro-4,6,6,7,8,8-hexamethylcyclopenta- $\gamma$ -2-benzopyran (CAS RN 1222-05-5)*. United States Environmental Protection Agency, Washington DC, USA. Accessed November 2020 at [https://www.epa.gov/sites/production/files/2015-09/documents/hhcb\\_wp\\_ra\\_final\\_08\\_27\\_14.pdf](https://www.epa.gov/sites/production/files/2015-09/documents/hhcb_wp_ra_final_08_27_14.pdf).

US EPA (2015). *Chemical and Product Categories Database (CPCat)*. United States Environmental Protection Agency, Washington DC, USA. Accessed December 2020 at <https://actor.epa.gov/cpcat/faces/search.xhtml>.

US EPA (2016). *2016 Chemical Data Reporting (CDR) Industrial Processing and Use Database*. United States Environmental Protection Agency, Washington DC, USA. Downloaded August 2020. Available at <https://www.epa.gov/chemical-data-reporting/using-2016-cdr-database>.

US EPA (2017a). *Estimations Programs Interface (EPI) Suite™ for Microsoft Windows® v 4.11*. United States Environmental Protection Agency (US EPA), Washington DC, USA. Available at <https://www.epa.gov/tsca-screening-tools/epi-suitetm-estimation-program-interface>.

US EPA (2017b). *The ECOSAR (ECOLOGical Structure Activity Relationship) Class Program for Microsoft Windows® v2.0*. United States Environmental Protection Agency, Washington DC, USA. Available at <https://www.epa.gov/tsca-screening-tools/ecological-structure-activity-relationships-ecosar-predictive-model>.

US EPA (2020a). *Chemical and Product Categories (CPCat) Chemical Use for 1,3,4,6,7,8-hexahydro-4,6,6,7,8,8-hexamethylindeno[5,6-c]pyran (CAS RN 1222-05-5)*. United States Environmental Protection Agency, Washington DC, USA. Downloaded November 2020. Available at <https://actor.epa.gov/cpcat/faces/chemicalUse.xhtml?casrn=1222-05-5>.

US EPA (2020b). *High Production Volume List*. United States Environmental Protection Agency Washington DC, USA. Accessed September 2020 at [https://comptox.epa.gov/dashboard/chemical\\_lists/EPAHPV](https://comptox.epa.gov/dashboard/chemical_lists/EPAHPV).

US EPA (2020c). *TSCA Chemical Substance Inventory*. United States Environmental Protection Agency Washington DC, USA. Accessed September 2020 at <https://www.epa.gov/tsca-inventory>.

Vallecillos L, Borrull F and Pocurull E (2014). On-line coupling of solid-phase extraction to gas chromatography–mass spectrometry to determine musk fragrances in wastewater. *Journal of Chromatography A*, **1364**, pp 1-11.

Villa S, Vighi M and Finizio A (2014). Theoretical and experimental evidences of medium range atmospheric transport processes of polycyclic musk fragrances. *Science of The Total Environment*, **481**, pp 27-34.

Wang L, McDonald JA and Khan SJ (2013). Enantiomeric analysis of polycyclic musks in water by chiral gas chromatography–tandem mass spectrometry. *Journal of Chromatography A*, **1303**, pp 66-75.

Wang L and Khan SJ (2014). Enantioselective analysis and fate of polycyclic musks in a water recycling plant in Sydney (Australia). *Water Science and Technology*, **69**(10), pp 1996-2003.

Wong F, Robson M, Melymuk L, Shunthirasingham C, Alexandrou N, Shoeib M, Luk E, Helm P, Diamond Miriam L and Hung H (2019). Urban sources of synthetic musk compounds to the environment. *Environmental Science: Processes & Impacts*, **21**(1), pp 74-88.

Xie Z, Ebinghaus R, Temme C, Heemken O and Ruck W (2007). Air–Sea Exchange Fluxes of Synthetic Polycyclic Musks in the North Sea and the Arctic. *Environmental Science & Technology*, **41**(16), pp 5654-5659.

Yamauchi R, Ishibashi H, Hirano M, Mori T, Kim J-W and Arizono K (2008). Effects of synthetic polycyclic musks on estrogen receptor, vitellogenin, pregnane X receptor, and cytochrome P450 3A gene expression in the livers of male medaka (*Oryzias latipes*). *Aquatic Toxicology*, **90**(4), pp 261-268.

