



Australian Government

Department of Health, Disability and Ageing

Australian Industrial Chemicals Introduction Scheme

C4 and C5 alkanolic acids

Evaluation statement (EVA00180)

26 June 2025



Table of contents

Contents

AICIS evaluation statement (EVA00180)	4
Subject of the evaluation.....	4
Chemicals in this evaluation.....	4
Reason for the evaluation	4
Parameters of evaluation	4
Summary of evaluation	5
Summary of introduction, use and end use.....	5
Human health.....	5
Proposed means for managing risk.....	7
Workers.....	7
Conclusions	8
Supporting information	9
Grouping rationale	9
Chemical identity	9
Relevant physical and chemical properties	11
Introduction and use	11
Australia	11
International	11
Existing Australian regulatory controls	12
AICIS.....	12
Public	12
Workers.....	12
International regulatory status.....	13
Exposure standards	13

Health hazard information	13
Toxicokinetics.....	13
Acute toxicity	14
Corrosion/Irritation.....	17
Sensitisation.....	20
Repeat dose toxicity	21
Genotoxicity	23
Carcinogenicity.....	25
Reproductive and development toxicity	25
References	29

AICIS evaluation statement (EVA00180)

Subject of the evaluation

C4 and C5 alkanolic acids

Chemicals in this evaluation

CAS name	CAS number
Propanoic acid, 2,2-dimethyl-	75-98-9
Propanoic acid, 2-methyl-	79-31-2
Pentanoic acid	109-52-4
Butanoic acid, 2-methyl-	116-53-0
Butanoic acid, 3-methyl-	503-74-2

Reason for the evaluation

Evaluation Selection Analysis indicated a potential human health risk.

Parameters of evaluation

The chemicals in this evaluation are branched and linear alkanolic acids containing 4 and 5 carbons that are listed on the Australian Inventory of Industrial Chemicals (the Inventory). These chemicals are grouped together based on their structural similarity and are expected to have similar hazard profiles.

This evaluation is a human health risk assessment for all identified industrial uses of these chemicals. Throughout the report, these chemicals will be referred to by their more common names, as follows:

- pivalic acid for propanoic acid, 2,2-dimethyl- (CAS No. 75-98-9)
- isobutyric acid for propanoic acid, 2-methyl- (CAS No. 79-31-2)
- valeric acid for pentanoic acid (CAS No. 109-52-4)
- 2-methylbutyric acid for butanoic acid, 2-methyl- (CAS No. 116-53-0)
- isovaleric acid for butanoic acid, 3-methyl- (CAS No. 503-74-2).

Summary of evaluation

Summary of introduction, use and end use

There is currently limited specific information about the introduction, use and end use of these chemicals in Australia. Pivalic acid has reported uses in personal care products (hair conditioner) in Australia (concentration unknown).

Based on international use information, these chemicals have functional use as pH regulating agents and fragrances. As such these chemicals may be used in a range of personal care products (cosmetics) and domestic products. Reported concentrations as fragrances in finished products are low with concentrations < 0.03% in most personal care and domestic product categories.

These chemicals have various site-limited applications with functional uses as intermediates and pH-regulators in the manufacture of chemicals and materials.

Some chemicals in this group have non-industrial uses, including as flavouring in food products.

Human health

Summary of health hazards

The identified health hazards are based on available data for chemicals in this group. In addition, alcohols are known to be metabolised to the respective acids. Therefore, read across information from relevant alcohols has been used to support the hazards associated with systemic exposure where appropriate.

Based on available data, these chemicals are expected to be readily absorbed following oral, dermal and inhalation exposure.

Based on the available data, these chemicals:

- have low acute oral, dermal and inhalation toxicity (with the exception of isobutyric acid)
- are not considered to be skin sensitisers
- are not expected to cause serious systemic health effects following repeated exposure
- are not expected to cause specific adverse effects on fertility/sexual function and foetal development
- are not expected to have genotoxic potential
- are not expected to be carcinogenic.

Isobutyric acid is currently classified as harmful if swallowed and in contact with skin. Although available data do not fully support these classifications sufficient data are not available to recommend an amendment.

Based on the available data, most of these chemicals are considered to be corrosive and cause severe damage to eyes. In dermal irritation studies in rabbits, these chemicals caused corrosive effects (visible necrosis) following exposures ≥ 3 minutes. In most eye irritation studies, the observed effects including corneal opacity and iritis were not reversible within the

study duration of 7–21 days. The available data for pivalic acid indicate that it is less irritating to skin and eyes but the severity of effects observed still warrant classification as a skin (mean erythema score > 2.3) and eye irritant (corneal opacity 2/4, iritis 1.87/2, conjunctival redness 2/3, chemosis 2.83/4).

For further details of the health hazard information see **Supporting information**.

Hazard classifications relevant for worker health and safety

These chemicals satisfy the criteria for classification according to the Globally Harmonized System of Classification and Labelling of Chemicals (GHS) (UNECE 2017) for hazard classes relevant for work health and safety as follows. This evaluation does not consider classification of physical hazards and environmental hazards.

Health hazards	Hazard category	Hazard statement
Acute toxicity	Acute Tox. 4	H302: Harmful if swallowed
Acute toxicity	Acute Tox. 4	H312: Harmful in contact with skin
Skin corrosion/irritation	Skin Corr. 1B	H314: Causes severe skin burns and eye damage
Serious eye damage/eye irritation	Eye Damage 1	H318: Causes serious eye damage
Skin corrosion/irritation	Skin Irrit. 2	H315: Causes skin irritation
Serious eye damage/eye irritation	Eye Irrit. 2	H319: Causes serious eye damage

For these hazard classifications, note that:

- The acute toxicity classifications apply only to propanoic acid, 2-methyl- (isobutyric acid; CAS No. 79-31-2). These classifications are currently listed in the Hazardous Chemical Information System.
- The corrosion and eye damage classifications apply to all chemicals except propanoic acid, 2,2-dimethyl- (pivalic acid; CAS No. 75-98-9).
- The skin and eye irritation classifications apply only to propanoic acid, 2,2-dimethyl- (pivalic acid; CAS No. 75-98-9).

Summary of health risk

Public

Based on the available use information, these chemicals may be used in a wide range of personal care products (cosmetic) and domestic products. The public may be exposed to these chemicals by:

- direct application of personal care products containing the chemicals to the skin, hair and lips
- direct skin contact during use of domestic products
- incidental skin and eye contact with the chemicals during use of domestic products
- inhaling aerosols/vapours.

The main route of exposure to these chemicals is expected to be via the skin. Incidental ingestion and eye contact may also occur. The critical health effect for risk characterisation of these chemicals is skin and eye corrosion.

Chemicals in this group function as pH regulating agents and fragrances. When used as a pH regulator, exposure to high concentrations of the free acid is not expected. Available information indicate use as a fragrance is only at low concentrations (< 0.03%). Therefore corrosive effects are not expected and; hence, there are no identified risks to the public that require management.

Workers

During product formulation and packaging, oral, dermal, inhalation and ocular exposure might occur, particularly where manual or open processes are used. These could include transfer and blending activities, quality control analysis, and cleaning and maintaining equipment. Worker exposure to these chemicals at lower concentrations could also occur while using formulated products containing these chemicals. The level and route of exposure will vary depending on the method of application and work practices employed. Good hygiene practices to minimise incidental oral exposure are expected to be in place.

Given the local health effects, these chemicals could pose a risk to workers. Control measures to minimise oral, dermal, ocular and inhalation exposure are needed to manage the risk to workers (see **Proposed means for managing risk** section).

Proposed means for managing risk

Workers

Recommendation to Safe Work Australia

It is recommended that Safe Work Australia (SWA) update the Hazardous Chemical Information System (HCIS) to include classifications relevant to work health and safety (see **Summary of health hazards** section).

Information relating to safe introduction and use

The information in this statement including recommended hazard classifications, should be used by a person conducting a business or undertaking at a workplace (such as an

employer) to determine the appropriate controls under the relevant jurisdiction Work Health and Safety laws.

Recommended control measures that could be implemented to manage the risk arising from oral, dermal, ocular and inhalation exposure to these chemicals include, but are not limited to:

- using closed systems or isolating operations
- minimising manual processes and work tasks through automating processes
- adopting work procedures that minimise splashes and spills
- cleaning equipment and work areas regularly
- using protective equipment that is designed, constructed, and operated to ensure that the worker does not come into contact with these chemicals.

Measures required to eliminate, or manage risk arising from storing, handling and using hazardous chemicals depend on the physical form and the manner in which these chemicals are used.

Personal protective equipment should not solely be relied upon to control risk and should only be used when all other reasonably practicable control measures do not eliminate or sufficiently minimise risk.

Model codes of practice, available from the Safe Work Australia website, provide information on how to manage the risks of hazardous chemicals in the workplace, prepare an SDS and label containers of hazardous chemicals. Your Work Health and Safety regulator should be contacted for information on Work Health and Safety laws and relevant Codes of Practice in your jurisdiction.

Conclusions

The Executive Director is satisfied that the identified risks to human health from the introduction and use of these industrial chemicals can be managed.

Note:


1. Obligations to report additional information about hazards under section 100 of the *Industrial Chemicals Act 2019* apply.
2. You should be aware of your obligations under environmental, workplace health and safety and poisons legislation as adopted by the relevant state or territory.


Supporting information

Grouping rationale


Chemicals in this evaluation consist of branched and linear alkanolic acids containing 4 or 5 carbon atoms. These chemicals are grouped together based on their similar structure, physicochemical properties and use profiles.

Chemical identity


CAS number	75-98-9
CAS name	Propanoic acid, 2,2-dimethyl-
Molecular formula	C ₅ H ₁₀ O ₂
Associated names	Neopentanoic acid Pivalic acid
Molecular weight (g/mol)	102.13
SMILES (canonical)	O=C(O)C(C)(C)C
Structural formula	

CAS number	79-31-2
CAS name	Propanoic acid, 2-methyl-
Molecular formula	C ₄ H ₈ O ₂
Associated names	Isobutyric acid
Molecular weight (g/mol)	88.11
SMILES (canonical)	O=C(O)C(C)C
Structural formula	


CAS number	109-52-4
CAS name	Pentanoic acid
Molecular formula	C ₅ H ₁₀ O ₂
Associated names	<i>n</i> -Valeric acid Propylacetic acid
Molecular weight (g/mol)	102.13
SMILES (canonical)	O=C(O)CCCC
Structural formula	



CAS number	116-53-0
CAS name	Butanoic acid, 2-methyl-
Molecular formula	C ₅ H ₁₀ O ₂
Associated names	2-Methylbutyric acid
Molecular weight (g/mol)	102.13
SMILES (canonical)	O=C(O)C(C)CC
Structural formula	



CAS number	503-74-2
CAS name	Butanoic acid, 3-methyl-
Molecular formula	C ₅ H ₁₀ O ₂
Associated names	3-Methylbutyric acid Isobutylformic acid Isopropylacetic acid

	Isovaleric acid
	Isovaleric acid
Molecular weight (g/mol)	102.13
SMILES (canonical)	O=C(O)CC(C)C
Structural formula	

Relevant physical and chemical properties

Chemicals in this group have molecular weights ranging between 88.11 and 102.13 g/mol. Most of these chemicals, with the exception of pivalic acid, are liquid at ambient temperatures, with boiling points between 154–187°C. Pivalic acid is solid at ambient temperatures with melting and boiling points at ~35°C and ~164°C, respectively. The calculated vapour pressures for these chemicals range from 95.59–436 Pa at 25°C (calculated with EPI MPBPVP), and the water solubility values from 15,590–49,180 mg/L (calculated with EPI WATERNT). The dissociation constant value (pK_a) for these chemicals is 4.9 at 20°C and log K_{ow} (calculated with EPI WSKOW) between 1–1.56 at 25°C (Danish QSAR n.d.).

Introduction and use

Australia

Limited specific information is available on the introduction, use and end use of these chemicals in Australia. Pivalic acid has identified use in personal care products (hair conditioner) on Australian retail websites (concentration unknown).

International

The following international uses have been identified through:

- Galleria Chemica (Chemwatch n.d.)
- Substances in Preparations in Nordic Countries (SPIN) database (SPIN n.d.)
- European Union (EU) Registration, Evaluation and Authorisation of Chemicals (REACH) dossiers
- the European Cosmetic Ingredient Database (CosIng) (EC n.d.)
- United States Environmental Protection Agency Chemical Data Reporting (CDR) (US EPA 2016; US EPA 2020)
- International Fragrance Association (IFRA) Transparency List
- RIFM Fragrance Ingredient Safety Assessments (Api et al. 2019a; Api et al. 2019b; Api et al. 2020)
- EWG's Skin Deep (EWG n.d.)

- Skinsort (Skinsort n.d.).

Chemicals in this group have reported use in personal care products (cosmetics) as perfuming and buffering agents.

2-Methylbutyric acid, isovaleric acid, isobutyric acid and pentanoic acid are on the IFRA transparency list. As fragrances they may be used in a wide range of personal care products and domestic products. Reported concentrations in finished products are low, with concentrations < 0.03% in most personal care and domestic product categories and < 0.5% in products not intended for direct skin contact such as scented candles (Api et al. 2019a; Api et al. 2019b; Api et al. 2020).

Chemicals in this group have reported site-limited applications with functional uses as intermediates and pH-regulators in the manufacture of chemicals and materials.

Some of these chemicals also have reported non-industrial uses as flavouring in food products.

Existing Australian regulatory controls

AICIS

No specific controls are currently available for these chemicals.

Public

No specific controls are currently available for these chemicals.

Workers

Two of these chemicals are listed in the Hazardous Chemical Information System (HCIS) (SWA n.d.) with the following hazard category and statements for human health:

Propanoic acid, 2-methyl- (isobutyric acid; CAS No. 79-31-2)

Health hazards	Hazard category	Hazard statement
Acute toxicity	Acute Tox. 4	H302: Harmful if swallowed
Acute toxicity	Acute Tox. 4	H312: Harmful in contact with skin

Pentanoic acid (valeric acid; CAS No. 109-52-4)

Health hazards	Hazard category	Hazard statement
Skin corrosion/irritation	Skin Corr. 1B	H314: Causes severe skin burns and eye damage

No exposure standards are available for these chemicals in Australia (SWA n.d.).

International regulatory status

Exposure standards

The following exposure standards were identified (Chemwatch n.d.):

- Time weighted average (TWA): 5 mg/m³ in Latvia and Russia (CAS No. 109-52-4); and 2 mg/m³ in Russia (CAS No. 503-74-2)
- Short-term Exposure Limit (STEL): 0.03 mg/m³ in Lithuania (CAS No. 79-31-2); 5 mg/m³ in Belarus (CAS No. 109-52-4); and 2 mg/m³ in Belarus (CAS No. 503-74-2).

Health hazard information

Health hazard data are available for 4 chemicals in this evaluation: pivalic acid, valeric acid, isovaleric acid and isobutyric acid. This information has been used to infer the toxicity of all chemicals in this evaluation in the absence of data.

In addition, alcohols are known to be metabolised to the respective acid by alcohol and aldehyde dehydrogenases (Government of Canada 2023; OECD 2003). Therefore, read across information from relevant alcohols has been used to support the hazard information associated with systemic exposure where appropriate.

Toxicokinetics

Based on the available data for isobutyric acid, isovaleric acid and relevant physicochemical properties, chemicals in this group are expected to be readily absorbed through the skin, lungs and gastrointestinal tract. These chemicals are expected to be rapidly metabolised, and the metabolites are excreted in urine, faeces or through exhaled air.

In a non-guideline study, Charles River CD rats were administered ¹⁴C-labelled isobutyric acid (4 males/dose) at 4, 40, and 400 mg/kg bw and 4 females at 400 mg/kg bw by gavage. The chemical was rapidly absorbed. The majority of the chemical was eliminated through respiration as ¹⁴CO₂, with 67–83% eliminated within 4 hours and 85–90% within 48 hours. Radioactivity excreted in urine and faeces averaged 3.5% and < 1%, respectively. In rats dosed with 400 mg/kg bw, isobutyric acid was rapidly cleared from the plasma, with levels peaking between 0.5 and 1 hour (11.4 ± 2.4 µg/mL), decreasing to 3.3 µg/mL at 1 hour, and were below the limit of detection at 4 hours (REACH n.d.-b).

With the exception of pivalic acid, the chemicals are expected to be metabolised via the fatty acid pathway or tricarboxylic cycle (JECFA 1999). In an *in vivo* toxicokinetic study, unlabelled isobutyric acid was administered to 10 male rats at a dose of 1 g/day in feed for 6 days. An increase in the urinary excretion levels of methylmalonic and succinic acids was observed (REACH n.d.-b).

In an *in vivo* toxicokinetic study, radiolabelled isovaleric acid administered to Sprague Dawley (SD) rats in feed was completely utilised in the production of cholesterol and fatty acids (Api et al. 2019b). Isovaleric acid is expected to undergo beta-oxidation and is metabolised to carbon dioxide through acetyl-CoA. It is also established that isovaleric acid is ketogenic, forming a carboxyl moiety and an isopropyl moiety, which can undergo cholesterol or fatty acid synthesis.

Most of the chemicals in this group are endogenous metabolites in humans, except for pivalic acid. These chemicals are not expected to bioaccumulate in the body.

Pivalic acid is reported to be metabolised primarily through esterification with carnitine, forming pivaloylcarnitine (Colnot and Dekant 2024).

Acute toxicity

Oral

Based on the weight of evidence, most chemicals in this group are considered to have low acute toxicity following oral exposure.

Isobutyric acid (CAS No. 79-31-2) is classified for Acute toxicity Category 4 (H302: harmful if swallowed). Although the available data from guideline studies do not support the current classification, given the effects seen in a dermal toxicity study there is not sufficient evidence to amend this classification.

Isobutyric acid

In a GLP-compliant acute oral toxicity study conducted in accordance with the Organisation for Economic Co-operation and Development (OECD) Test Guideline (TG) 401, WISW (SPF THO) rats (5/sex/dose) were treated with isobutyric acid at 1990, 2510, 2835 or 3160 mg/kg bw via gavage. The reported median lethal dose (LD50) was 2230 mg/kg bw with some deaths observed at all doses. Reported signs of toxicity included sedation, ataxia, ruffled fur, hypothermia and staggering (doses not specified) (REACH n.d.-b).

In a non-GLP compliant acute oral toxicity study similar to OECD TG 401, Gassner rats (5/sex/dose) were treated with isobutyric acid in water at 200, 1600, 2500, 2800, 3200 or 6400 µL/kg bw (equivalent to 190, 1516, 2370, 2650, 3030 or 6060 mg/kg bw) via gavage. The reported LD50 was 2510 mg/kg bw. Reported signs of toxicity included abdominal position, apathy, dyspnoea and cyanosis (doses not specified) (REACH n.d.-b).

Pivalic acid

In a non-GLP compliant acute oral toxicity study similar to OECD TG 420, male SD rats (5/dose) were treated with pivalic acid in corn oil at 34.6, 120, 417, 1450, 5000 or 10000 mg/kg bw via gavage. The reported LD50 was 2000 mg/kg bw. Reported signs of toxicity included depressed activity, severe dyspnoea, depressed reflexes, sprawling, incoordination, and abnormal gait. These were observed at doses at which lethality was observed ≥ 1450 mg/kg bw (REACH n.d.-a).

Valeric acid

In a GLP compliant acute oral toxicity study conducted in accordance with OECD TG 401, female Wistar rats (5/dose) were treated with valeric acid in water at 1250, 1600 or 2000 mg/kg bw via gavage. The reported LD50 was 1720 mg/kg bw. Reported signs of toxicity (dose not specified) included crouching, shrunken flanks, piloerection, reduced activity, abdominal lateral position, high stepping and unsteady gait, irregular respiration and absence of rightening reflex (REACH n.d.-c).

In a non-GLP compliant acute oral toxicity study similar to OECD TG 401, SD rats (5/sex/dose) were treated with valeric acid in olive oil at 2150, 3160, 4640 or 6810 mg/kg bw

via gavage. The reported LD50 was 4600 mg/kg bw. Reported signs of toxicity (dose not specified) included irregular respiration, apathy, abdominal position, staggering, narcosis-like state, absent pain reflex, piloerection, diarrhoea, exsiccosis (insufficient intake of fluids) and bloody excrements (REACH n.d.-c).

Isovaleric acid

In an acute oral toxicity study similar to OECD TG 401 (GLP status not specified), SD rats (5/sex/dose) were treated with isovaleric acid in carboxymethyl cellulose at 681, 1000, 1470, 2150, 2610, 3160, 3830 or 5000 mg/kg bw via gavage. The reported LD50 was 2500 mg/kg bw. Signs of toxicity including dyspnoea, apathy, unkempt fur, and poor condition were reported at doses \geq 1000 mg/kg bw, cyanosis, atony, unsteady gait at doses \geq 1470 mg/kg bw, and narcosis, impaired pain reflexes, trembling, exsiccation and exophthalmia at 2610 mg/kg bw (REACH n.d.-e).

Oral LD50 values below 2000 mg/kg bw/day were reported (CCOHS 2024a; CCOHS 2024b; CCOHS 2024c; CCOHS 2024d; CCOHS 2024e). However, no information on test methodologies was available.

Dermal

Based on the available data, most chemicals in this group are considered to have low acute toxicity following dermal exposure.

Isobutyric acid (CAS No. 79-31-2) has a classification Acute toxicity Category 4 (H312: Harmful in contact with skin). Although the limited available data indicate an LD50 might be below 1000 mg/kg bw/day, given limitations of this study, the available data do not support an amendment to the current classification.

Isobutyric acid

In a non-GLP compliant non-guideline acute dermal toxicity study, male New Zealand White (NZW) rabbits (4/dose) were treated with graduated single doses of isobutyric acid. The number and quantity of individual doses were not specified. The study had some limitations including limited reporting and the use of occlusive dressing. The reported LD50 was 0.5 mL/kg bw (equivalent to 474 mg/kg bw) with a range of 0.37 to 0.67 mL/kg bw (equivalent to 351 to 636 mg/kg bw) (REACH n.d.-b).

Pivalic acid

In a non-GLP compliant acute dermal toxicity study similar to OECD TG 402, NZW rabbits (4/sex/dose) were treated with a single dose of pivalic acid at 50, 200, 794 or 3160 mg/kg bw under occlusive conditions. The reported dermal LD50 was 3160 mg/kg bw. Reported clinical signs of toxicity at doses at which lethality occurred (3160 mg/kg bw) included depression, severe dyspnoea, prostration, excessive urination and coma (REACH n.d.-a).

Valeric acid

In a non-GLP compliant acute dermal toxicity study similar to OECD TG 402, Wistar rats (6/sex/dose) were treated with a single dose of valeric acid (unknown vehicle) at 2000 or 4000 mg/kg bw under occlusive conditions over 24 hours. The reported dermal LD50 was > 2000 mg/kg bw. Reported clinical signs of toxicity (doses not specified) included irregular

breathing, giddiness, atonia, narcotic-like behaviour, insensibility to pain, dry and rough coat, diarrhea and bloody stool (REACH n.d.-c).

Isovaleric acid

In an acute dermal toxicity study similar to OECD TG 402, Vienna White rabbits (3/sex) were treated with a single dose of isovaleric acid in olive oil at 2000 mg/kg bw under semi-occlusive conditions over 24 hours. The reported dermal LD50 was > 2000 mg/kg bw. No clinical signs of toxicity were reported (REACH n.d.-e).

Dermal LD50 values below 2000 mg/kg bw/day were reported (CCOHS 2024a; CCOHS 2024b; CCOHS 2024d; CCOHS 2024e) however no information on test methodologies was available.

Inhalation

Based on the limited available data, chemicals in this group are considered to have low acute toxicity following inhalation exposure. Although the concentration at which studies were conducted does not enable a definitive comparison with classification criteria to rule out acute toxicity, the absence of mortality in most studies indicate low toxicity.

Pivalic acid

In a GLP-compliant acute inhalation toxicity study conducted in accordance with OECD TG 436, Wistar rats (3/sex) were exposed to pivalic acid as an aerosol, with a mass median aerodynamic diameter (MMAD) of up to 0.26 µm, nose-only for 4 hours at a mean concentration of 5.30 mg/L. The reported median lethal concentration (LC50) for rats was > 5.30 mg/L. Reported clinical signs of toxicity included hunched posture, piloerection and wet fur (REACH n.d.-a).

In a non-GLP compliant acute inhalation toxicity study similar to OECD TG 403, Swiss mice, Wistar rats and Hartley guinea pigs (10/sex) were exposed to pivalic acid as a vapour to whole body for 6 hours at a nominal concentration of 4 mg/L. Two rats and one guinea pig died within 2 days post exposure indicating an LC50 for rats and guinea pigs of > 4 mg/L. All mice died within 24 hours of exposure (REACH n.d.-a).

Isobutyric acid

In a non-GLP compliant acute inhalation toxicity study similar to OECD TG 403, rats (strains unspecified) (6/sex) were exposed to isobutyric acid as a vapour for 8 hours at a concentration of 9.59 mg/L. The LC50 value could not be determined from this study as all animals survived. Dyspnoea was reported in one animal (REACH n.d.-b).

In a non-GLP compliant acute inhalation toxicity study similar to OECD TG 403, albino rats (6/sex) were exposed to isobutyric acid as a vapour, whole body for 8 hours at a concentration of 7.2 mg/L. The LC50 value could not be determined from this study as all animals survived. No clinical signs were reported in this study (REACH n.d.-b).

Valeric acid

In a non-guideline acute inhalation toxicity study, SD rats (6/sex) were treated with valeric acid as vapour, whole body for 7 hours at a concentration of 11.63 mg/L. The LC50 value

could not be determined from this study as all animals survived. No clinical signs were reported in this study (REACH n.d.-c).

In an acute inhalation toxicity study similar to OECD TG 403, rats (6/sex, strains unspecified) were exposed to isovaleric acid as vapour (type of inhalation exposure not specified) for 7 hours at a concentration of 2.06 mg/L (584 ppm). The LC50 value could not be determined from this study as all animals survived. No clinical signs were reported in this study (REACH n.d.-e).

Specific target organ toxicity – single exposure

Narcotic effects

Based on the effects observed in acute toxicity studies summarised above, the chemicals in this group are considered to cause transient narcotic effects after acute exposures.

Signs of narcosis were consistently observed for all chemicals in acute oral and dermal toxicity studies (see **Acute Toxicity** section) These included:

- decreased activity or lethargy
- lack of coordination
- ataxia (loss of muscle control).

However, where data are available, it indicates that these effects occurred at dose levels associated with mortality or approaching the lethal dose. Based on the weight of evidence, classification is not warranted.

Corrosion/Irritation

Skin irritation

Based on the available data, most chemicals in this group cause skin corrosion warranting classification.

Valeric acid (CAS No. 109-52-4) has a classification for Skin corrosion Category 1B (H314: Causes severe skin burns and eye damage). The available data support the current classification. Although necrotic effects were observed following 3 minute exposures it is not possible to determine whether these were observed within 1 hour. Therefore, sub-categorisation 1B is supported based on corrosive effects observed following 1 hour of exposure. This sub-categorisation is also appropriate for isovaleric acid. Based on observed effects in a non-guideline study and read across from valeric and isovaleric acid, hazard classification as a corrosive is recommended for isobutyric acid and 2-methylbutyric acid. The available data for pivalic acid indicate that pivalic acid is less irritating to skin but the severity of effects observed (erythema > 2.3) still warrant classification as a skin irritant.

Pivalic acid

In a GLP-compliant skin irritation study similar to OECD TG 404, undiluted pivalic acid (0.5 mL, assumed 100% purity) was applied to the skin of 6 NZW rabbits (sex not specified) for 4 hours under semi-occlusive conditions. Observations were recorded at 24, 48 and 72 hours, and at 7 days after patch removal. The following mean scores were reported for observations at 24, 48 and 72 hours: 3.61 for erythema and 1.83 for oedema, respectively (maximum

score of 4). Individual animal scores were not reported. Skin irritation effects were not reversible in all animals within 7 days (REACH n.d.-a).

In a non-GLP compliant skin irritation study similar to OECD TG 404, the skin of NZW rabbits (4/sex) were treated with 0.5 mL of pivalic acid (purity not reported) for 24 hours under occlusive (intact and abraded) conditions. Observations were recorded at 24 and 72 hours, and at 7 days after patch removal. The following mean scores were reported for observations at 24 and 72 hours: 2.8 for erythema and 2.55 for oedema, respectively (maximum score of 4). Individual animal scores were not reported. Skin irritation effects were not reversible in all animals within 7 days (REACH n.d.-a).

Isobutyric acid

Undiluted isobutyric acid (99% purity) was applied to 2 Vienna White rabbits (sex unspecified) under occlusive conditions for 20 hours. Observations were recorded at 24, 48 and 72 hours, and at 8 days. The following mean scores for individual animals were recorded based on observations at 24, 48 and 72 hours: 2.33, 2 for erythema and 3, 2.66 for oedema, respectively (maximum score of 4). Necrosis was observed in all animals and the effects were not reversible within 8 days (REACH n.d.-b).

Valeric acid

In a GLP compliant skin irritation study conducted in accordance with OECD TG 404, the skin of 3 NZW rabbits (sex unspecified) was applied with undiluted valeric acid (0.5 mL, 99.2% purity) for one hour under semi-occlusive conditions. Observations were recorded at 1, 24, 48 and 72 hours, and at 7 and 14 days after patch removal. The following mean scores for individual animals were recorded based on observations at 24, 48 and 72 hours: 4*, 2, 4* for erythema and 0.33, 1.66, 1.33 for oedema, respectively (maximum score of 4). Maximum scores of 4, as signified by an asterisk, were provided where scoring was not possible due to severe hardening, bulging or discolouring of the skin. Severe skin lesions and scar formation were observed in all animals and the effects were not reversible within 14 days (REACH n.d.-c).

In a non-guideline study, undiluted valeric acid (0.5 mL, purity not reported) was applied to the shaved skin of 2 Vienna White rabbits (sex unspecified) for 3 minutes or 4 hours. Observations were recorded at 24, 48, 72 hours and 8 days. The following mean scores for individual animals were recorded based on observations at 24, 48 and 72 hours: 4, 4 for erythema and 2, 2 for oedema for 3 minute exposure, and 4, 4 for erythema and 2, 2 for oedema for 4 hour exposure, respectively (maximum score of 4). Full thickness necrosis was observed in all animals and the effects were not reversible within 8 days (REACH n.d.-c).

Application of 0.01 mL of undiluted chemical (reported as a mixture of valeric acid and 2-methylbutyric acid, 99.4% active ingredient) to shaved skin of NZW rabbits resulted in moderate to severe irritation. The effects were not reversible within 14 days (REACH n.d.-c). No other details are available for this study.

Isovaleric acid

In a skin irritation study similar to OECD TG 404, rabbits (strain unspecified, n = 4 and 2 for 3 minute and 1 hour exposure, respectively) were treated with undiluted isovaleric acid (purity not reported) and observed for 8 days. The following mean scores for individual animals were recorded based on observations at 24 and 48 hours: 4, 2, 4, 2 for erythema and 2, 2, 2, 2.5 for oedema for 3 minute exposure, and 4, 3 for erythema and 2, 2 for oedema for 1 hour exposure, respectively (maximum score of 4). Full thickness necrosis was observed in all

animals after 1 hour exposure and the effects were not reversible within 8 days (REACH n.d.-e).

In another skin irritation study similar to OECD TG 404, Vienna White rabbits (2 males and 4 females) were treated with 0.5 mL of undiluted isovaleric acid for 4 hours under occlusive conditions. Observations were made for 24, 48 hours and 8 days after patch removal. The following mean scores were recorded based on observations at 24 and 48 hours: 4 for erythema and 1.33 for oedema. Individual animal scores were not reported. Necrosis was observed in all animals and the effects were not reversible within 8 days (REACH n.d.-e).

Eye irritation

Based on the available data, most of the chemicals in this group are expected to cause serious eye damage, warranting classification.

Isobutyric acid, valeric acid and isovaleric acid are corrosive to skin. Chemicals that are corrosive to skin are deemed to cause serious eye damage. This is supported by irreversible eye effects including corneal opacity and iritis observed in available studies for isobutyric acid and valeric acid. The available data for pivalic acid indicate that pivalic acid is less irritating to eyes but the severity of effects observed still warrant classification as an eye irritant. No data are available for 2-methylbutyric acid. In the absence of data and based on structural similarity, read across from isobutyric acid, valeric acid and isovaleric acid is considered appropriate.

Pivalic acid

In a non-GLP compliant eye irritation study similar to OECD TG 405, 3 mg of pivalic acid was instilled into the eyes of NZW rabbits (6 replicates). Observations were recorded at 1, 4, 24, 48 and 72 hours, and at 7 days. The following mean scores were reported based on observations at 24, 48 and 72 hours: corneal opacity 0.11/4, iritis 0.11/2, conjunctival redness 1.7/3. Individual animal scores were not reported. Irritation effects were reversible in all animals within 4 days (REACH n.d.-a).

In a non-GLP compliant eye irritation study similar to OECD TG 405, 0.2 mL of pivalic acid (undiluted) was instilled into the conjunctival sac of both eyes for each of 2 NZW rabbits. Observations were recorded at 1–2, 24, 48 and 72 hours, and at 7 days. The following mean scores were reported based on observations at 24, 48 and 72 hours: corneal opacity 2/4, iritis 1.87/2, conjunctival redness 2/3, chemosis 2.83/4. Individual animal scores were not reported. Irritation effects were not fully reversed within 7 days but showed signs of healing (REACH n.d.-a).

Isobutyric acid

In a non-guideline eye irritation study (BASF standard), 0.05 mL of isobutyric acid (undiluted, 99% purity) was instilled into the conjunctival sac of one eye for each of 2 Vienna White rabbits. Observations were recorded at 24, 48 hours and at 8 days. The following mean scores were reported based on observations at 24 and 48 hours: corneal opacity 3/4, iritis 0/2, conjunctival redness 0.3/3, chemosis 2/4. Iritis developed over time in both animals with a mean score of 2/2 at day 8. Individual animal scores were not reported. Irritation effects were not reversible within 8 days (REACH n.d.-b).

Valeric acid

In a non-guideline eye irritation study (BASF standard), 0.05 mL of valeric acid (undiluted) was instilled into the conjunctival sac of one eye for each of 3 White Vienna rabbits. Observations were recorded at 24, 48, 96 hours and at 8 days. The following mean scores for individual animals were reported based on observations at 24, 48 and 96 hours:

- animal 1, corneal opacity 2.33/4, iritis 0.66/2, conjunctival redness 2/3, chemosis 2/4
- animal 2, corneal opacity 2.66/4, iritis 1/2, conjunctival redness 2/3, chemosis 1.67/4
- animal 3, corneal opacity 2.33/4, iritis 1/2, conjunctival redness 2/3, chemosis 2.33/4 (REACH n.d.-c).

Effects showed no signs of reversibility by day 8 (termination of study).

In a GLP-compliant eye irritation study conducted in accordance with OECD TG 405, 0.01 mL of undiluted chemical (reported as a mixture valeric acid and 2-methylbutyric acid, 99.4% active ingredient) was instilled into the conjunctival sac of both eyes of a male NZW rabbit. Observations were recorded at 1, 24, 48 and 72 hours and at 7, 10, 14, 17 and 21 days. The following mean scores were reported based on observations at 24, 48 and 72 hours: corneal opacity 4/4, iritis 1.33/2, conjunctival redness 3/3, chemosis 2.33/4. Severe ocular injury, including corneal erosion and bulging were observed and the effects were not reversible within 21 days (REACH n.d.-c).

Observation in humans

In a human patch test study, 1% isovaleric acid in petrolatum was not irritating to the skin following 48 hours of occlusive exposure (REACH n.d.-e). No further study details were available.

Sensitisation

Skin sensitisation

Limited data are available for chemicals in this group. Based on the available data, the chemicals are not considered to be skin sensitisers.

In a non-GLP compliant guinea pig maximisation test (GPMT) similar to OECD TG 406, guinea pigs (10/sex, strain unspecified) received intradermal injection of 0.05% (w/v) pivalic acid in corn oil and topical application of 25% (w/v). The animals were challenged with topical application of the chemical at 10% (w/v) in corn oil under occlusive conditions and dermal reactions were observed at 24 and 48 hours. No reactions were observed following challenge. The chemical was not considered to be a skin sensitiser (REACH n.d.-a).

Observation in humans

In 2 human maximisation tests, 2-methylbutyric acid (Api et al. 2019a) and isovaleric acid (1% in petrolatum) (Api et al. 2019b; REACH n.d.-e) were found not to have caused skin sensitisation. No further study details were available.

In silico

These chemicals have no structural alerts for protein binding based on the mechanistic profiling functionality of the Organisation for Economic Co-operation and Development

(OECD) QSAR Application Toolbox (OECD QSAR Toolbox v4.5) (OECD 2020). These chemicals are predicted to be non-sensitising using OASIS–TIMES (Optimised Approach based on Structural Indices Set–Tissue Metabolism Simulator; version 2.31) (OASIS LMC), and the knowledge-based expert system Deductive Estimation of Risk from Existing Knowledge (DEREK) Nexus version 6.0.1 (Lhasa Limited).

Repeat dose toxicity

Oral

Based on the limited data available, chemicals in this group are not expected to cause serious systemic health effects following repeated oral exposure.

Pivalic acid

In a GLP-compliant repeated dose toxicity study conducted similar to OECD TG 407, Fischer 344 rats (7/sex/dose) were administered pivalic acid in water/polyethylene glycol (PEG) 200 (50/50 v/v) by gavage at 10, 30, 100 or 300 mg/kg bw/day daily for 28 days. No treatment-related mortality was reported. A mild irritation effect (dark nasal discharge) was observed in animals immediately after dosing with the chemical at 100 and 300 mg/kg bw/day. Changes (direction and magnitude not reported) in alkaline phosphatase, cholesterol, creatinine and bilirubin were observed in females at 100 and 300 mg/kg bw/day, indicating a change in liver function. Minor changes in hepatic macroscopic appearance, liver weight and kidney weight were also reported. However, there were no associated histopathological changes. The no observed adverse effect level (NOAEL) was determined to be 30 mg/kg bw/day for both sexes (REACH n.d.-a).

In a GLP-compliant repeated dose toxicity study conducted in accordance with OECD TG 408, SD rats (10/sex/dose) were administered pivalic acid in 1% methylcellulose with 0.1% Tween 80 by gavage at 25, 125 or 300–375 mg/kg bw/day daily for 90 days. The highest dose level was reduced from 375 to 300 mg/kg bw/day from Day 23 due to unexpected deaths. Four rats from the highest dose group were euthanised due to deterioration in general condition and breathing impairment. Macroscopic examination showed disturbance of the gastrointestinal tract (presence of gas and abnormal coloured faecal content) as a result of the irritant nature of the chemical. No other treatment-related mortality occurred during the study. Reported clinical signs of toxicity in the highest dose group included wet and/or dry rales, piloerection, hunched posture, decreased activity and body weight loss. No significant treatment-related effects were observed in ophthalmoscopic examination, haematology, urinalysis, clinical chemistry, organ weights, gross pathology or histopathology parameters at any dose level compared to control animals. The established NOAEL in the study was 125 mg/kg bw/day for both sexes (REACH n.d.-a).

Isobutyric acid

In a subchronic repeated dose toxicity study similar to OECD TG 408, Crj: CD(SD) rats (30/sex/dose) were administered isobutyric acid in water by gavage at 100, 316 or 1000 mg/kg bw/day daily for 92 days. Mortality was observed in control and dosed groups (1/60, 1/60, 2/60 and 11/60 for the control, 100, 316 and 1000 mg/kg bw/day, respectively) although it was reported to be not related to the systemic toxicity of the chemical. In the highest dose group, reported clinical signs of toxicity included hypoactivity, ataxia and salivation. No significant treatment-related effects were observed in ophthalmoscopic examination, haematology, urinalysis, clinical chemistry, organ weights, gross pathology or histopathology parameters at any dose level compared to control animals. The no observed

effect level (NOEL) value was determined to be 316 mg/kg bw/day for both sexes (REACH n.d.-b).

Isovaleric acid

In a non-guideline repeated dose toxicity study, neutralised isovaleric acid (isovaleric acid content: 40% w/w) was administered to 6 male SD rats at 5% (50,000 ppm, equivalent to 2500 mg/kg bw/day) in feed for 90 days. No mortality was reported. No significant treatment-related effects were reported in food consumption, bodyweight development, organ weights, haematology, blood chemistry, urinalysis, or histopathology parameters (Api et al. 2019b; REACH n.d.-e). The NOAEL value was determined to be 2500 mg/kg bw/day.

Structurally related alcohols – read across

No adverse effects up to the highest dose tested were observed in repeated dose oral toxicity studies with 1-pentanol (CAS No. 71-41-0), isobutanol (CAS No. 78-83-1) and isoamyl alcohol (CAS No. 123-1-3) (Api et al. 2025; Government of Canada 2023; NICNAS 2013). This included:

- a 13 week oral gavage study in rats (1-pentanol and isobutanol)
- a combined repeated dose toxicity study with the reproduction/developmental toxicity screening test (isoamyl alcohol)
- a 13 week drinking water study in rats (isoamyl alcohol)
- a 17 week oral gavage study in rats (isoamyl alcohol).

Dermal

Limited data are available for systemic health effects following repeated dermal exposure. Local effects consistent with skin corrosion were observed.

Pivalic acid

In a non-guideline repeated dose toxicity study, the skin of rabbits (strain unspecified) (4/sex/dose) was treated with undiluted pivalic acid at 30 or 300 mg/kg bw/day under occlusive conditions, once daily for 10 days. Slight to moderate erythema, atonia and desquamation was observed in both groups. Gross pathological findings showed parasitic cysts in the liver and fibrous, pitted kidneys, congestion of the lungs and the pancreas. No NOAEL value was established in the study (REACH n.d.-a).

Valeric acid

In a non-guideline 14 day repeated dose toxicity study, NZW rabbits (5/sex) were administered valeric acid (25% w/w in mineral oil) by dermal application at an equivalent of 500 mg/kg bw/day, 5 days/week for 2 weeks. One treated animal died on day 7. Severe erythema, moderate to severe oedema, necrosis and eschar formation, desquamation (peeling) and fissuring of the skin, and exfoliation of eschar tissue was observed in the application sites of treated animals. Histological findings were related to the skin irritation of the chemical. Organ weights, haematological, clinical, and urinary parameters were not examined in the study. No NOAEL value was established in the study (REACH n.d.-c).

In a non-guideline combined chronic repeated dose and carcinogenicity study, 50 male C3H/HeJ mice were administered undiluted valeric acid (purity not specified) by dermal application at 25 mg/kg bw/day (reduced from 50 mg/kg bw/day from week 3) 2 days/week

for 80 weeks. Mean body weight was lower in the treated group (83%) compared to the control group. Mortality was reported in the treated group; however, it was considered to be related to dermal toxicity of the chemical. Skin reactions (ulceration, scar formation) were observed at this dose. Tubular necrosis was observed in kidneys of 3 out of 19 animals examined (REACH n.d.-c). No NOAEL value for repeated dose toxicity was established in the study.

Inhalation

No data are available for the chemicals. Based on the read-across data available for isobutanol, chemicals in this group are not expected to cause serious systemic health effects following repeated inhalation exposure.

Structurally related alcohols – read across

No adverse effects up to the highest dose tested were observed in repeated dose inhalation toxicity studies with isobutanol (CAS No. 78-83-1) (Government of Canada 2023; NICNAS 2013).

This included:

- a 13 week inhalation study in rats
- a 2 generation reproductive toxicity study in rats (whole-body inhalation exposure).

Genotoxicity

Based on the available data, chemicals in this group are not expected to be genotoxic. The chemicals were mostly negative in *in vitro* studies. Although valeric acid was positive in an *in vitro* chromosome aberration study, there are no indications of genotoxicity *in vivo*.

In vitro

The available *in vitro* data for these chemicals were reported as mostly negative.

Pivalic acid

Negative results were reported in:

- a bacterial reverse mutation assay (Ames test) (OECD TG 471) in *Salmonella typhimurium* strains TA 98, TA 100, TA 1535 and TA 1537, and *Escherichia coli* WP2 uvrA with and without metabolic activation (S9) at concentrations of 50–5000 µg/plate (REACH n.d.-a)
- a mammalian chromosome aberration assay (OECD TG 473) in Chinese hamster ovary (CHO) cells with and without metabolic activation at concentrations up to 1020 µg/mL (REACH n.d.-a)
- a mammalian cell gene mutation study (OECD TG 476) in the thymidine kinase (TK) locus in mouse lymphoma cells L5178Y with and without metabolic activation at concentrations up to 1021 µg/mL (REACH n.d.-a).

Isobutyric acid

Negative results were reported in:

- 4 bacterial reverse mutation assays (OECD TG 471) in *S. typhimurium* strains TA 98, TA 100, TA 1535, TA 1537 and TA 1538, and *E. coli* WP2 uvrA with and without metabolic activation at concentrations up to 5000 µg/plate (Api et al. 2020; REACH n.d.-b)
- a mammalian cell gene mutation study (OECD TG 476) in the hypoxanthine-guanine phosphoribosyl transferase (HPRT) locus in CHO cells with and without metabolic activation at concentrations up to 440 µg/mL (REACH n.d.-b)
- a micronucleus test (OECD TG 487) in human lymphocytes with and without metabolic activation at concentrations up to 880 µg/mL (Api et al. 2020).

Valeric acid

The following results were reported:

- negative in a bacterial reverse mutation assay (OECD TG 471) in *S. typhimurium* strains TA 98, TA 100, TA 1535, TA 1537 and TA 1538 with and without metabolic activation at concentrations of 10,000 µg/plate (REACH n.d.-c)
- positive in a mammalian chromosome aberration assay (OECD TG 473) in CHO cells with and without metabolic activation at concentrations up to 1200 µg/mL (REACH n.d.-c)
- negative without metabolic activation and positive with metabolic activation in a sister chromatid exchange (SCE) assay (OECD TG 479) in CHO cells at concentrations up to 2000 µg/mL (REACH n.d.-c)
- negative in a mammalian cell gene mutation study (OECD TG 476) in the HPRT locus in CHO cells with and without metabolic activation at concentrations up to 1000 µg/mL (cytotoxic limit) (REACH n.d.-c).

2-Methylbutyric acid

Negative results were reported in:

- a bacterial reverse mutation assay (OECD TG 471) in *S. typhimurium* strains TA 97a, TA 98, TA 100, TA 102 and TA 1535 with and without metabolic activation at concentrations of 5000 µg/plate (Api et al. 2019a).

Isovaleric acid

Negative results were reported in:

- two bacterial reverse mutation assays (OECD TG 471) in *S. typhimurium* strains TA 98, TA 100, TA 102, TA 1535 and TA 1537, and *E. coli* WP2 uvrA with and without metabolic activation at concentrations up to 5000 µg/plate (Api et al. 2019b; REACH n.d.-e).

In vivo

Valeric acid

In a GLP compliant mammalian erythrocyte micronucleus test similar to OECD TG 474, Swiss Webster mice (5/sex/dose) were treated with valeric acid by intraperitoneal injection at single doses within the range between 50–400 mg/kg bw (equivalent to 25%, 50% and 80% of the average LD50 value of 332 mg/kg bw for males and females) in corn oil. The incidence

of micronuclei in peripheral blood polychromatic erythrocytes did not increase significantly in any of the treated groups, indicating a lack of clastogenicity (REACH n.d.-c).

Structurally related alcohols – read across

Isobutanol and isoamyl alcohol were negative in an *in vivo* mammalian erythrocyte micronucleus test conducted in accordance with OECD TG 474 with NMRI mice. The incidence of micronuclei in bone marrow polychromatic erythrocytes did not increase significantly in any of the treated groups, indicating a lack of clastogenicity (Api et al 2025; Government of Canada 2023; NICNAS 2013).

In silico

Chemicals in this group present no structural alerts for *in vivo* mutagenicity based on molecular structure as profiled by the OECD QSAR Toolbox v4.5 (OECD 2020). The expert rule based system, SARAH Nexus version 3.1.0 (Lhasa Limited n.d.) identified the alerting group (mutagenicity *in vitro*) for pivalic acid, but no structural alert presented for the rest of the group. These chemicals were predicted to be *in vitro* Ames negative in OASIS TIMES (OASIS LMC).

Carcinogenicity

Limited data are available for chemicals in this group. There is low concern for carcinogenicity of these chemicals as they are non-genotoxic and did not cause effects in repeated dose toxicity studies. The equivocal evidence of carcinogenicity for valeric acid in a dermal study appears related to the corrosive nature of the chemical.

In a non-guideline combined chronic repeated dose and carcinogenicity study (see **Repeat dose toxicity – dermal** section), incidences of squamous cell carcinomas (4 mice), fibrosarcomas (6 mice), fibromas (3 mice), and keratoacanthoma (1 mouse) were reported in male mice administered valeric acid by dermal application at 25 mg/kg bw/day (reduced from 50 mg/kg bw/day from week 3) 2 days/week for 80 weeks. Histopathological re-examination revealed that the observed neoplasms (with the exception of 3 tumours) were associated with scar tissue formation and skin repair mechanisms, rather than from an inherent carcinogenic potential of the chemical (REACH n.d.-c).

In silico

Chemicals in this group have no alerts for carcinogenicity (genotoxic and non-genotoxic) based on the molecular structure as profiled by the OECD QSAR Toolbox v4.5 (OECD 2020).

Reproductive and development toxicity

Based on the available data, these chemicals are not expected to cause specific adverse effects on sexual function or foetal development. Observed developmental effects are considered secondary to maternal toxicity.

Pivalic acid

In a GLP compliant prenatal developmental toxicity study conducted in accordance with OECD TG 414, pregnant SD rats (20/dose) were administered pivalic acid (vehicle: 1% methylcellulose with 0.1% Tween 80) by gavage once daily at 25, 75 or 225 mg/kg bw/day

on gestational days (GD) 6–19. Dams were sacrificed on GD 20 and the developmental and reproductive toxicity parameters were examined. No treatment-related mortality was reported. Clinical signs of toxicity were observed at the highest dose including wet/dry rales and piloerection. No significant changes in the level of thyroid hormones (T3, T4 and TSH) were reported. An increase in incidence of post-implantation losses, as a result of higher early and late resorptions was reported at the highest dose. The mean foetal weights (male, female and combined) were significantly lower in this group compared to controls. No significant changes in sex ratio, anogenital distance (AGD), the mean numbers of corpora lutea and implantations or pre-implantation losses were reported in all treated groups. Nipple retention was not examined in the study. At 225 mg/kg bw/day, skeletal (partially split sternum) and visceral (heart and major vessel) abnormalities were reported. Skeletal variations (bent scapula, delayed ossification) and an increase in incidence of enlarged isthmus/absent lobe of thyroid is considered to be associated with low body weight and foetal immaturity, and not an adverse effect of the chemical. The reported NOAEL value was 75 mg/kg bw/day for both maternal toxicity and developmental toxicity (REACH n.d.-a).

In a GLP-compliant extended one generation reproductive toxicity study conducted in accordance with OECD TF 443, SD rats (F0: 25/sex/dose; F1: 20/sex/dose) were administered pivalic acid (vehicle: 1% methylcellulose with 0.1% Tween 80) by gavage daily at 15, 50 and 100 (F0: 100/150) mg/kg bw/day including 10 weeks prior to mating for F0 parents. Treatment-related mortality was reported in F0 and F1 generations at 50 or 100 mg/kg bw/day, and it is considered to be associated with irritant effect of the chemical rather than systemic toxicity. Reported clinical signs of toxicity included abnormal breathing and distended abdomen. No adverse effects on body weight, food consumption, clinical pathology, thyroid hormones, macro- and micro-pathology were reported in either the F0 or F1 generation. Reproductive performance for the F0 adults, litter size, offspring survival and bodyweight were unaffected by parental treatment at the highest dose. No treatment-related effect on sexual maturation, ovarian follicle and corpora lutea counts were reported. Increased thyroid and kidney weights (with no microscopic correlations) and higher T4 serum levels (with no effect on TSH levels) were observed in both generations. No significant changes in sex ratio, AGD or nipple retention were reported in all dosed groups. The reported NOAEL value was 100 mg/kg bw/day for both systemic toxicity and reproductive/developmental toxicity (REACH n.d.-a).

In a GLP compliant prenatal developmental toxicity study conducted in accordance with OECD TG 414, pregnant NZW rabbits (24/dose) were administered pivalic acid (vehicle: carboxymethyl cellulose (CMC)) by gavage daily at 12.5, 25 or 50 mg/kg bw/day on GD 6–28. Mortality was reported in the 25 and 50 mg/kg bw/day groups (3 animals in each group). The gravid uterine weight for females treated with 50 mg/kg bw/day was significantly lower than controls, consistent with the lower litter weight observed at this dose. No significant changes in the number of resorptions, implantation losses, number of foetuses or sex ratio were reported in all treated groups. Mean foetal and litter weights were significantly lower in the highest dose group compared to controls. However, it is not considered to be associated with any adverse effect on *in utero* survival or morphological development. The reported NOAEL values were determined to be 12.5 mg/kg bw/day for maternal toxicity and 50 mg/kg bw/day for developmental toxicity (REACH n.d.-a).

Valeric acid

In a prenatal developmental toxicity study similar to OECD TG 414, 22 pregnant SD rats were administered valeric acid (> 99%) in water by gavage daily at 750 mg/kg bw/day on GD 6–15. Mortality was reported in the treated group (2 out of 22 animals). Clinical signs of toxicity included wheezing, dyspnoea, salivation, rough coat and hunched position. Gross pathological changes in stomach and kidney were reported in 10 out of 22 animals. There

were no significant changes in maternal and foetal body weight, food consumption, the number of corpora lutea, implantations, resorptions (early and late) or sex ratio. Domed head was observed in one foetus (out of 160) in the treated group. No treatment-related external, skeletal and visceral malformations were reported. Slight increase in incidences of skeletal variation (delayed ossification) was considered to be associated with maternal toxicity in the tested group. The lowest observed adverse effect level (LOAEL) was determined to be 750 mg/kg bw/day (only dose tested) for maternal toxicity based on clinical signs. The reported NOAEL for reproductive and developmental toxicity was 750 mg/kg bw/day (REACH n.d.-c).

In a non-guideline study described as an *in vivo* developmental toxicity screen and a conventional Segment II-type protocol (US EPA), pregnant SD rats (20–24/dose) were administered valeric acid (> 99%) in corn oil by gavage daily on GD 6–15. In the screening test, animals received 75 or 1000 mg/kg bw/day, the dams were allowed to deliver, and their litters were examined through postnatal day 6. In the Segment II study, animals received 50, 100 or 200 mg/kg bw/day and caesarean sections were performed on GD 20. Mortality was reported in all treated groups (1, 3 and 10 out of 24 animals at 50, 100 and 200 mg/kg bw/day, respectively). Clinical signs of toxicity including rales and dyspnoea were observed in all treated groups. Reduced maternal weight gain was reported in all treated groups. There were no differences in number of corpora lutea, number of implantations, pre-implantation loss, post-implantation loss, or the number of live foetuses per litter in all treated groups compared to controls. Reduced foetal body weight was reported at 100 and 200 mg/kg bw/day and sternebral variations at 50, 100 and 200 mg/kg bw/day. The study reported the foetal effects were as a result of maternal toxicity (REACH n.d.-c).

A further re-analysis of the data generated from the study above revealed that the sternebral variations (delayed ossification and offset sternebrae) lacked a dose related trend, were transient, and were present in all groups including controls. The reported NOAEL value for developmental toxicity was 50 mg/kg bw/day due to decreased foetal body weight. No NOAEL value for maternal toxicity was determined due to toxicity effects at all tested doses (REACH n.d.-c).

Isovaleric acid

In a GLP compliant prenatal developmental toxicity study similar to OECD TG 414, 10 pregnant Wistar rats were administered isovaleric acid (99.9%) in olive oil by gavage once daily at 600 mg/kg bw/day on GD 6–19. Dams were sacrificed on GD 20 and the developmental and reproductive toxicity parameters were examined. No treatment-related mortality was reported. Salivation and local irritation of the larynx and upper and lower respiratory tract were reported in treated dams, but these effects were not considered to be severe. There were no significant changes in the number of resorptions, implantation losses, number of foetuses or sex ratio. No increased in incidences of external, skeletal or visceral malformations or variations were observed in the treated group. The reported NOAEL for maternal and developmental toxicity was 600 mg/kg bw/day (only dose tested) (Api et al. 2019b; REACH n.d.-e).

Structurally related alcohols – read across

In reproductive and developmental toxicity studies with 1-pentanol (CAS No. 71-41-0), isobutanol (CAS No. 78-83-1) and isoamyl alcohol (CAS No. 123-1-3), no adverse effects on fertility and sexual function were observed and any developmental effects (decreased foetal weight and minor skeletal variation) were considered secondary to maternal toxicity (Api et al. 2025; Government of Canada 2023; NICNAS 2013). This included:

- OECD TG 416 2 generation reproductive toxicity study (whole-body inhalation) in rats (isobutanol)
- OECD TG 422 combined repeated dose toxicity study with the reproduction/developmental toxicity screening test (isoamyl alcohol)
- a whole body inhalation developmental study (1-pentanol)
- an OECD 443 Extended One-Generation Reproductive Toxicity study (isoamyl alcohol)
- numerous OECD 414 prenatal development studies (isobutanol and isoamyl alcohol).

References

- Api AM, Belsito D, Biserta S, Botelho D, Bruze M, et al. (2019a) '[RIFM fragrance ingredient safety assessment, 2-methylbutyric acid, CAS Registry Number 116-53-0](#)', *Food and Chemical Toxicology*, 130: 110574, DOI: 10.1016/j.fct.2019.110574.
- Api AM, Belsito D, Biserta S, Botelho D, Bruze M, et al. (2019b) '[RIFM fragrance ingredient safety assessment, isovaleric acid, CAS Registry Number 503-74-2](#)', *Food and Chemical Toxicology*, 130: 110570, DOI: 10.1016/j.fct.2019.110570.
- Api AM, Belsito D, Biserta S, Botelho D, Bruze M, et al. (2020) '[RIFM fragrance ingredient safety assessment, isobutyric acid, CAS Registry Number 79-31-2](#)', *Food and Chemical Toxicology*, 144: 111673, DOI: 10.1016/j.fct.2020.111673.
- Api AM, Bartlett A, Belsito D, Botelho D, Bruze M, et al. (2025) '[Update to RIFM fragrance ingredient safety assessment, isoamyl alcohol, CAS Registry Number 123-51-3](#)', *Food and Chemical Toxicology*, 197: 115240, DOI: 10.1016/j.fct.2025.115240.
- CCOHS (Canadian Centre for Occupational Health and Safety) (2024a) [Record for propanoic acid, 2,2-dimethyl-, CAS No. 75-98-9](#), CCOHS, accessed 21 November 2024.
- CCOHS (Canadian Centre for Occupational Health and Safety) (2024b) [Record for propanoic acid, 2-methyl-, CAS No. 79-31-2](#), CCOHS, accessed 21 November 2024.
- CCOHS (Canadian Centre for Occupational Health and Safety) (2024c) [Record for pentanoic acid, CAS No. 109-52-4, CCOHS](#), accessed 21 November 2024.
- CCOHS (Canadian Centre for Occupational Health and Safety) (2024d) [Record for butanoic acid, 2-methyl-, CAS No. 116-53-0](#), CCOHS, accessed 21 November 2024.
- CCOHS (Canadian Centre for Occupational Health and Safety) (2024e) [Record for butanoic acid, 3-methyl-, CAS No. 503-74-2](#), CCOHS, accessed 21 November 2024.
- Colnot T and Dekant W (2024) 'Grouping of short alkyl-chain branched carboxylic acids for developmental toxicity', *Regulatory Toxicology and Pharmacology*, 151: 105662, DOI: 10.1016/j.yrtph.105662
- Chemwatch (n.d.) [Galleria Chemica](#), Chemwatch website, accessed 21 November 2024.
- Danish QSAR (Danish Environmental Protection Agency) (n.d.) [Danish QSAR](#), accessed 9 January 2025.
- EC (European Commission) (n.d.) [CosIng](#), EC website, accessed 25 November 2024.
- EWG (n.d.) [EWG's Skin Deep](#), EWG website, accessed 5 February 2025.
- Government of Canada (2023) [Draft screening assessment - Selected C3-C5 Alcohols Group](#), Government of Canada, accessed 27 March 2025.
- IFRA (International Fragrance Association) (n.d.) [Transparency List](#), IFRA website, accessed 21 November 2024.

JECFA (Joint FAO/WHO Expert Committee on Food Additives) (1999). Evaluation of certain food additives and contaminants. Forty-ninth report of the JointFAO/WHO Expert Committee on Food Additives. Rome, 17-26 June 1997. WHO Technical ReportSeries, no. 884. Geneva.

Lhasa Limited DEREK Nexus (version 6.0.1), [Computer software], Lhasa Limited, accessed 9 January 2025.

Lhasa Limited SARAH Nexus (version 3.1.0), [Computer software], Lhasa Limited, accessed 9 January 2025.

NICNAS (National Industrial Chemicals Notification and Assessment Scheme) (2013) [*IMAP Single Assessment Report – 1-Propanol, 2-methyl-, CAS Number 78-83-1: Human health tier II assessment*](#), NICNAS, accessed 21 November 2024.

OASIS LMC (Laboratory of Mathematical Chemistry) Optimised Approach based on Structural Indices Set–Tissue Metabolism Simulator (OASIS–TIMES) (Version 2.31), [Computer software], LMC, accessed 9 January 2025.

OECD (The Organisation for Economic Co-operation and Development) (2003) [*SIDS Initial Assessment Profile, SIAM 17*](#), OECD, accessed 21 November 2024.

OECD (Organisation for Economic Co-operation and Development) (2020) [*Quantitative Structure-Activity Relationship Toolbox \(Version 4.5\)*](#) [Computer software], OECD website, accessed 9 January 2025.

REACHa (Registration, Evaluation, Authorisation and Restriction of Chemicals) (n.d.) [*Registered dossier for propanoic acid, 2,2-dimethyl-, CAS No. 75-98-9*](#), European Chemicals Agency website, accessed 21 November 2024.

REACHb (Registration, Evaluation, Authorisation and Restriction of Chemicals) (n.d.) [*Registered dossier for propanoic acid, 2-methyl-, CAS No. 79-31-2*](#), European Chemicals Agency website, accessed 21 November 2024.

REACHc (Registration, Evaluation, Authorisation and Restriction of Chemicals) (n.d.) [*Registered dossier for pentanoic acid, CAS No. 109-52-4*](#), European Chemicals Agency website, accessed 21 November 2024.

REACHd (Registration, Evaluation, Authorisation and Restriction of Chemicals) (n.d.) [*Registered dossier for butanoic acid, 2-methyl-, CAS No. 116-53-0*](#), European Chemicals Agency website, accessed 21 November 2024.

REACHe (Registration, Evaluation, Authorisation and Restriction of Chemicals) (n.d.) [*Registered dossier for butanoic acid, 3-methyl-, CAS No. 503-74-2*](#), European Chemicals Agency website, accessed 21 November 2024.

Skinsort (n.d.) [*SkinSort*](#), Skinsort website, accessed 5 February 2025.

SPIN (Substances in Preparation in Nordic Countries) (n.d.) [*SPIN Database*](#), SPIN website, accessed 21 November 2024.

SWA (Safe Work Australia) (n.d.) [*Hazardous Chemicals Information System*](#), SWA website, accessed 21 November 2024.

TGA (Therapeutic Goods Administration) (2024) [Therapeutic Goods \(Poisons Standard—June 2024\) Instrument 2024](#), TGA, accessed 4 December 2024.

UNECE (United Nations Economic Commission for Europe) (2017) [Globally Harmonized System of Classification and Labelling of Chemicals \(GHS\), Seventh Revised Edition](#), UNECE, accessed 4 December 2024.

US EPA (United States Environmental Protection Agency) (2016) [Access Chemical Data Reporting Data \(CDR data\)](#), US EPA website, accessed 4 December 2024.

US EPA (United States Environmental Protection Agency) (2020), [Access Chemical Data Reporting Data \(CDR data\)](#), US EPA website, accessed 4 December 2024.

