

Existing Chemical Hazard Assessment Report



Australian Government
Department of Health and Ageing
NICNAS

Dimethyl Terephthalate

June 2008

© Commonwealth of Australia 2008

This work is copyright. You may download, display, print and reproduce this material in unaltered form only (retaining this notice) for your personal, non-commercial use or use within your organisation. Apart from any use as permitted under the Copyright Act 1968, all other rights are reserved. Requests and inquiries concerning reproduction and rights should be addressed to Commonwealth Copyright Administration, Attorney General's Department, Robert Garran Offices, National Circuit, Barton ACT 2600 or posted at <http://www.ag.gov.au/cca>

Preface

This report was compiled under the National Industrial Chemicals Notification and Assessment Scheme (NICNAS). This Scheme was established by the *Industrial Chemicals (Notification and Assessment) Act 1989* (Cwlth) (the Act), which came into operation on 17 July 1990.

The principal aim of NICNAS is to aid in the protection of people at work, the public and the environment from the harmful effects of industrial chemicals.

NICNAS assessments are carried out in conjunction with the Department of the Environment, Water, Heritage and the Arts, which carry out the environmental assessment for NICNAS. NICNAS has two major programs: the assessment of the health and environmental effects of new industrial chemicals prior to importation or manufacture; and the other focussing on the assessment of chemicals already in use in Australia in response to specific concerns about their health/or environmental effects.

There is an established mechanism within NICNAS for prioritising and assessing the many thousands of existing chemicals in use in Australia.

For the purposes of Section 78(1) of the Act, copies of assessment reports for New and Existing Chemical assessments are freely available from the web (www.nicnas.gov.au). Summary Reports are published in the *Commonwealth Chemical Gazette* (<http://www.nicnas.gov.au/publications/#gazette>), and are available to the public on line at www.nicnas.gov.au.

Copies of this report and other NICNAS reports are available on the NICNAS website. Hardcopies are available from NICNAS at the following address:

**GPO Box 58
Sydney NSW 2001
AUSTRALIA
Attention: Office Manager
Tel: +61 (02) 8577 8800
Freecall: 1800 638 528
Fax: +61 (02) 8577 8888
Email: info@nicnas.gov.au**

Other information about NICNAS (also available on request) includes:

- NICNAS Annual Reports.
- NICNAS Service Charter.
- Brochure on NICNAS Registration.

More information on NICNAS can be found at the NICNAS web site:

<http://www.nicnas.gov.au>

Overview

This review of dimethyl terephthalate (DMT) is a health hazard assessment only. For this assessment, an OECD SIDS Initial Assessment Report on Dimethyl Terephthalate (DMT) was consulted. Information from this report was supplemented with literature surveys conducted up to September 2006.

DMT is used widely as an industrial intermediate to manufacture polyethylene terephthalate (PET) and dioctyl terephthalate.

In Australia, DMT is imported for distribution to various institutions and laboratories for analytical, pharmaceutical and biotechnological research.

Structurally, phthalate esters are characterized by a diester structure consisting of a benzenedicarboxylic acid head group linked to two ester side chains. However, there are three isomeric forms of benzenedicarboxylic acid esters: *ortho*-, *meta*- and *para*-phthalates, also known as phthalates, isophthalates and terephthalates, respectively. Terephthalate esters such as DMT have two carbon atoms between carboxylic acid functional groups and ester side chains. DMT is therefore structurally distinct from *ortho*-phthalate (commonly known simply as phthalate) esters.

Toxicity data for DMT were available for the majority of health endpoints. Additional data from the major metabolite terephthalic acid (TPA) were also available.

DMT was readily absorbed after oral administration. Elimination was rapid with urine being the major route of excretion. There was no evidence of accumulation in tissues after multiple doses. The hydrolysis product, TPA, was the only metabolite detected in the urine in rats, while urinary metabolites in mice consisted of monomethyl terephthalate (70%) and TPA (30%).

DMT has low acute oral, dermal and inhalational toxicity. DMT causes minimal skin and eye irritant effects in animals, and did not induce skin sensitisation in guinea pigs.

DMT does not appear to be mutagenic or genotoxic, nor was it deemed to be carcinogenic based on a 2-year feeding study.

For repeat dose toxicity, a conservative NOAEL based on reduced body weight gains in a rat 96-day dietary study was determined at 313 mg/kg bw/d. The target organ identified in oral exposure studies was the urinary tract and the subsequent formation of bladder crystals or calculi.

Following 90-days of exposure via the inhalation route, a NOAEL of 86.4 mg/m³ was determined. At this dose, no urinary effects were observed and clinical effects such as nose rubbing, preening, and blinking, were considered mild and transient.

In a one-generation reproduction study, DMT in the diet up to 1% (636 mg/kg bw/d) had no effect on reproduction although pup weights were reduced at weaning. There was no evidence of fertility effects, and hence the NOAEL for fertility was 636 mg/kg bw/d, the highest dose tested. In the same study, the NOAEL for developmental effects was 152 mg/kg bw/d, with a LOAEL of 313 mg/kg bw/d based on reduced pup weights at weaning, which was probably a lactational effect.

The NOAEC in an inhalation developmental study for DMT was 1 mg/m³ and NOAEL in an oral developmental study was 1000 mg/kg.

The DMT metabolite, TPA, was also examined in reproduction studies. In a one-generation reproduction feeding study, no adverse effects on fertility were noted in adult rats fed up to 5% TPA (2480-3018 mg/kg bw/d). While the NOAEL for fertility effects was 5.0% (2480-3018 mg/kg bw/d), the NOAEL for maternal toxicity was 0.5%. The NOAEL for developmental toxicity was 0.5% (240-307 mg/kg bw/d), based on increased postnatal deaths and decreased pup survivability at higher doses. The adverse effects observed in the offspring in this study appear to be the result of maternal toxicity and the formation of renal and bladder calculi in the weanling animals.

In a 2-generation study, rats treated with TPA showed no effects on various fertility indices including sperm motility, number and morphology. No overt changes in the developmental landmarks were observed. Anogenital distance was not decreased in male pups. Reductions in bodyweight generally occurred when offspring started to consume solid diet, and were considered to be a direct effect of the test material on the pups rather than related to developmental toxicity. The NOAEL for effects on both fertility and development was 20000 ppm, (2011-5072 mg/kg bw/d) the highest dose tested in the study.

In an inhalation development study, the NOAEC for TPA was 10 mg/m³.

In contrast, a single study showed no effects on serum testosterone or testes weight but significant decreases in sperm head counts and daily sperm production in rats fed 5% TPA in the diet (5818 mg/kg bw/d). In addition, several sperm motility parameters also reduced significantly in all treated groups including at the lowest dose of 0.2%. Reproductive performance was not measured in this study.

With respect to fertility effects, a more recent OECD test guideline 2-generation reproduction study of di-2-ethylhexyl terephthalate (DEHT), also known to be metabolised to TPA in rats, showed no effects on fertility indices in both generations up to the highest dose tested of 1% (F1: 893-1549 mg/kg bw/d) in the diet.

Therefore, overall, weight-of-evidence from available studies indicates that DMT or its metabolite TPA do not affect fertility, at doses up to approximately 5000 mg/kg bw/d. In addition, no developmental effects unrelated to maternal toxicity were observed with DMT in oral and inhalation studies.

Table of Contents

PREFACE	iii
OVERVIEW	iv
ACRONYMS AND ABBREVIATIONS	vii
1. INTRODUCTION	1
2. IDENTITY	2
2.1 Identification of the substance	2
2.2 Physicochemical properties	2
3. USES	3
4. HUMAN HEALTH HAZARD	4
4.1 Toxicokinetics	4
4.2 Acute toxicity	4
4.3 Irritation	5
4.3.1 Skin irritation	5
4.3.2 Eye irritation	5
4.3.3 Respiratory irritation	6
4.4 Sensitisation	6
4.5 Repeated dose toxicity	6
4.6 Genetic toxicity	9
4.7 Carcinogenicity	11
4.8 Reproductive toxicity	11
4.8.1 Repeat dose toxicity studies	12
4.8.2 Prenatal developmental toxicity studies	12
4.8.3 One/two generation reproductive toxicity studies	12
5. HAZARD CHARACTERISATION	15
6. HUMAN HEALTH HAZARD SUMMARY TABLE	17
REFERENCES	19

Acronyms and Abbreviations

bw	body weight
C	Celsius
CAS	Chemical Abstracts Service
CHO	Chinese hamster ovary
d	day
DMT	dimethyl terephthalate
DMSO	dimethyl sulphoxide
DNA	deoxyribonucleic acid
F1	filial 1 (first generation)
g	gram
GD	gestation day
GLP	good laboratory practice
h	hour
kg	kilogram
kPa	kilopascals
L	litre
LC50	median lethal concentration
LD50	median lethal dose
LOAEL	lowest-observed-adverse-effect level
mg	milligram
mL	millilitre
NICNAS	National Industrial Chemicals Notification and Assessment Scheme
NOAEC	no-observed-adverse-effect concentration
NOAEL	no-observed-adverse-effect level
OECD	Organisation for Economic Cooperation and Development
ppm	parts per million
SIDS	Screening Information Data Set
TPA	terephthalic acid
μ	micro

1. Introduction

This review of dimethyl terephthalate (DMT) is a health hazard assessment only. For this assessment, an OECD SIDS Initial Assessment Report on Dimethyl Terephthalate (DMT) (OECD, 2001) was consulted. Information from this report was supplemented with relevant studies from more recent literature surveys conducted up to September 2006.



Information on Australian uses was compiled from data supplied by industry in 2004 and 2006.

References not marked with an asterisk were examined for the purposes of this assessment. References not examined but quoted from the key report as secondary citations are also noted in this assessment and marked with an asterisk.

2. Identity

2.1 Identification of the substance

CAS Number: 120-61-6
 Chemical Name: 1,4-Benzenedicarboxylic acid, dimethyl ester
 Common Name: Dimethyl terephthalate (DMT)
 Molecular Formula: $C_{10}H_{10}O_4$
 Structural Formula:



Molecular Weight: 194.2
 Synonyms: Dimethyl 1,4-benzenedicarboxylate;
 Dimethyl p-benzenedicarboxylate;
 Dimethyl p-phthalate;
 Methyl 4-carbomethoxybenzoate;
 Methyl p-(methoxycarbonyl)benzoate;
 Terephthalic acid, dimethyl ester
 Purity/Impurities/Additives: Purity: 99.9% min.
 Impurity: 40 ppm max. methyl (p-formyl)benzoate;
 225 ppm max. methyl hydrogen terephthalate
 Additives: none

2.2 Physicochemical properties

Table 1: Summary of physicochemical properties

Property	Value
Physical state	White solid or colourless molten liquid
Melting point	141°C
Boiling point	Not available
Density	Not available
Vapour pressure	$<1.33 \times 10^{-3}$ kPa (25°C)
Water solubility	0.019 g/L (25°C)
Partition coefficient n-octanol/water (log Kow)	2.25
Henry's law constant	Not available
Flash point	153°C

Source: OECD (2001)

3. Uses

DMT is used widely as an industrial intermediate to manufacture polyethylene terephthalate (PET) and dioctyl terephthalate (OECD, 2001).

In Australia, DMT is imported for distribution to various institutions and laboratories for analytical, pharmaceutical and biotechnological research.

4. Human Health Hazard

4.1 Toxicokinetics

Previous evaluations

Oral

Following a single or repeated oral dose to rats (20 or 40 mg/kg bw), ^{14}C -DMT was readily absorbed from the digestive tract. Excretion was almost complete within 48 hours, with 75-81% in urine and 4-16% in faeces. There was no evidence of tissue accumulation after multiple doses (Moffitt, 1975). In rats, terephthalic acid (TPA) was the only metabolite detected in the urine suggesting that the hydrolysis of DMT to TPA was virtually complete, while urinary metabolites in mice consisted of monomethyl terephthalate (70%) and TPA (30%) (Heck, 1980*).

Dermal

When ^{14}C -DMT was applied to unabraded depilated rat skin (80 mg in 0.2 mL vehicle) either as a single dose or on alternate days for 10 days under occlusion, around 11% of a single dose and 13% of five repeated doses were excreted in the urine and faeces of rats within 10 days (Moffitt, 1975).

Ocular

When ^{14}C -DMT was applied to rabbit eye, approximately 29% and 37% of the administered dose (50 mg) were excreted in the urine following 5 min or 24 h exposure with no evidence of tissue accumulation or ocular damage. Faecal excretion was minimal ($\leq 2\%$) (Moffitt, 1975).

Data not reported in previous evaluations

No data.

Conclusion

Several studies conducted in rats and mice have indicated that DMT is readily absorbed from the digestive tract and rapidly eliminated in the urine within 48 hours. Most of absorbed DMT is metabolised to TPA via hydrolysis. In mice, the predominant metabolite was monomethyl terephthalate. DMT is also absorbed via dermal and ocular routes in rats and rabbits.

4.2 Acute toxicity

Previous evaluations

The results for acute oral, dermal and inhalation toxicity in laboratory animals are summarised in the Table below.

Study	Species	Results	References
Oral	Rat	LD50 = 4390 mg/kg bw LD50 > 6590 mg/kg bw	Marhold, 1972 * Krasavage et al., 1973
Dermal	Guinea pig	LD50 > 5000 mg/kg bw	Patty's Ind Hyg Tox, 1981*
Inhalation	Rat	LC50 > 6 mg/L	Sanina & Kochetkova, 1963*

Data not reported in previous evaluations

No data.

Conclusion

DMT has low acute oral, dermal and inhalation toxicity in laboratory animals.

4.3 Irritation

4.3.1 Skin irritation

Previous evaluations

Several unpublished studies were identified but lack details. DMT applied dermally to guinea pigs, mice (tails dipped into DMT solutions), and rabbits were reported to be slightly irritating (Eastman Kodak, 1957*, 1963*; Sanina & Kocketkova, 1963*).

Data not reported in previous evaluations

No data.

Conclusion

DMT caused minimal skin irritation in laboratory animals.

4.3.2 Eye irritation

Previous evaluations

In two studies, DMT instilled in rabbit eyes had either no effect or induced a mild irritation (Eastman Kodak, 1957*; Anonymous, 1986*). A third study indicated that a pronounced conjunctivitis was induced (time for return to normal was not reported) (Kamal'Dinova, 1962*).

Data not reported in previous evaluations

No data.

Conclusion

DMT caused minimal eye irritation in rabbits.

4.3.3 Respiratory irritation

Previous evaluations

No data.

Data not reported in previous evaluations

No data.

Conclusion

No respiratory irritation studies were available for assessment.

4.4 Sensitisation

Previous evaluations

One study was available for assessing the skin sensitisation potential of DMT (Krasavage et al., 1973). In this study, 0.5 mL of a 1% solution of DMT in acetone:dioxane:guinea pig fat (7:2:1) was placed onto the rump area of 10 animals. Primary irritation was determined at 24 and 48 hours post-dosing. After one week, this dosing solution was mixed with whole heparinized rabbit blood (1%) for 1 to 3 hours and 0.05 mL of this mixture was injected into the footpad of the animals. A challenge dose (0.5 mL of a 1% solution of DMT in acetone:dioxane:guinea pig fat (7:2:1) was administered one week later. Sensitisation scores were recorded at 24 and 48-hours after the challenge dose. Neither primary irritation nor skin sensitization was induced by DMT.

Data not reported in previous evaluations

No data.

Conclusion

DMT did not induce skin sensitisation in guinea pigs.

4.5 Repeated dose toxicity

Previous evaluations

Several repeat dose toxicity studies have been conducted via oral (gavage and dietary) and inhalation routes of exposure. Studies range in duration from 2-13 weeks for the oral route and up to 6 months for the inhalation route.

Bladder stone induction was investigated in weanling F344 rats (13-18/sex/dose group) fed DMT diets for a period of two weeks at 0, 0.5, 1.0, 1.5, 2, or 3% (average dose for males/females was 660/638, 1320/1277, 1890/1790, 2260/2290 and 2590/3020 mg/kg bw/d). Bodyweights were decreased at 1.5% and above in females and at 1% and above in males on days 6-8 and 12-14. Decreased bodyweights were accompanied by reduced food consumption. The incidence of bladder calculi in males from the 0, 0.5, 1.0, 1.5, 2.0 and 3.0% groups were 0, 0, 0, 35, 72 and 100%, respectively, and in females 0, 0, 0, 0, 36, and 47%, respectively. Irregular thickening of the bladder wall was limited to animals having bladder calculi. The composition of

the bladder calculi from the DMT treated animals was primarily calcium and TPA. The NOAEL was 0.5% (660 mg/kg bw/d for males) and 1.0% (1277 mg/kg bw/d for females) (Chin et al., 1981), based on bodyweight changes at higher doses.

Rats (7-19/group) were fed DMT via diet at 0.5, 1.6 or 3% for 13-weeks (Vogin, 1972*, cited in Heck & Tyl, 1985). The incidence of bladder calculi at 3% DMT was 12/16 and 6/16 in male and female rats, respectively. The incidence of bladder calculi in male rats was 1/19 at 1.6% and 2/19 at 0.5% DMT. Calculi were not noted in mid- and low-dose females. The incidence of mild to moderate hyperplasia of the bladder urothelium at 3% was 11/16 and 7/16 in males and females, respectively. Of the animals fed with 3% DMT diet and developing hyperplasia of the bladder lining, calculi were found in 11/11 males and 6/7 females. No NOAEL was determined.

Groups (10/sex/dose) of rats and mice were fed DMT in their diets at 1750, 2500, 5000, 10000, or 20000 ppm for 13-weeks (NCI, 1979*). No treatment-related effects were noted in physical appearance, behaviour or food consumption in either species. No deaths occurred in the rats. One male mouse at 2500, 5000, and 20000 ppm and two females at 20000 ppm died during the study. Reduced bodyweight gains were observed in both sexes of rats whereas body weight gains in mice were unaffected. No gross lesions were observed in either species at necropsy. Microscopic examination of the livers from both species in all dose groups revealed diffuse hepatocellular swelling but not in a dose-related manner. A NOAEL of 5000 ppm for rats and 20000 ppm for mice were derived.

DMT was evaluated in F344 rats and B6C3F1 mice (50/sex/dose) for carcinogenic potential in a 2-year dietary study (NCI, 1979*) (see Section 3.7). There were no effects on body weight or survival or clinical signs of toxicity reported from DMT in the diet at doses up to 0.5% (approximately 300 mg/kg bw/d), the highest dose tested.

Weanling male Long-Evans Hooded rats (30/group/dose) were exposed to DMT in the diet at 0, 0.25, 0.5 or 1% for 96 days (approximately 0, 152, 313 or 636 mg/kg bw/d) (Krasavage et al., 1973). Reduced weight gain was observed in the 1% dose animals. No effects were observed in haematological or biochemical parameters or absolute and relative liver and kidney weights. Microscope examinations were unremarkable. A NOAEL of 313 mg/kg bw/d was determined based on reduced weight gains at 636 mg/kg bw/d.

In a parallel study, male Long-Evans Hooded rats (30/dose) were exposed (whole body) to an atmosphere containing DMT at 0, 16.5, and 86.4 mg/m³ for 4-hours per day for a total of 58 exposures over a 3-month period (Krasavage et al., 1973). The particulate respirable fraction was 36%. Average daily doses were calculated at 0.7 and 4.0 mg/kg bw/d. Nose rubbing, preening and blinking were noted soon after the start of DMT exposures at the highest concentration. These symptoms continued intermittently throughout the exposure period and were repeated during succeeding exposures, but were not seen at the lower concentration. No toxicological effects were observed in biochemical or haematological parameters at any dose. Average and relative liver and kidney weights were unaffected. Tissue microscopic examination revealed no substance related changes. A NOAEC was assigned as 86.4 mg/m³, the highest dose tested.

Rats and guinea pigs were exposed to DMT at 15 mg/m³ for 6-hours per day, 5 days a week over a 6-month period (Lewis, 1982*). The respirable proportion was reported as 5 mg/m³. There were no effects on body weights, clinical chemistry or urinalysis parameters. Gross and histopathological evaluations were unremarkable. The NOAEC was 15 mg/m³.

Rats exposed to 0.08, 0.4 or 1 mg/m³ DMT by inhalation (described as a “chronic” exposure but duration not stated) induced a significant decrease (20% at 0.4 mg/m³, 51% at 1 mg/m³) in the uptake of radiolabelled noradrenaline by grey matter synaptosomes (Davidenko, 1984*). No effects were noted on monoamine oxidase or catecholamine-o-methyl transferase activity. No other details were provided.

Data not reported in previous evaluations

In a two-generation study conducted to OECD Test Guideline 416 and GLP, Wistar rats (26/group/sex) were fed diet containing 0, 1000, 5000 or 20000 ppm TPA (approximately 98 – 261, 492 – 1218, 2011 – 5072 mg/kg bw/d) (Milburn, 2003). The experimental design is described in Section 4.8.3 – One/Two generation reproductive toxicity studies.

Bodyweights were reduced in F0 and F1 males receiving 20000 ppm. Food consumption of F1 males and females receiving 20000 ppm was generally reduced and food utilization was less efficient in animals receiving this dose compared to controls in both generations.

Changes in the urinary bladder (multiple masses, firm deposits, blood stained or discoloured urine, thickening of the bladder wall and/or prominent blood vessels) indicating irritant effect on the bladder mucosa were observed in both sexes in the 20000 ppm group in both generations. The incidence was greater in F1 than in F0 animals. In all treated male groups, kidney weights (both absolute and relative) were decreased. However, kidney weight reductions were not accompanied by histological changes and were therefore regarded as physiological adaptation (Jacobsen, 2005). At 20000 ppm, minimal to slight renal papillary necrosis was reported in macroscopically abnormal kidneys of two F0 and 2 F1 males. Relative liver weights were also increased in both sexes and both generations at the highest dose level.

Conclusion

Collectively, data suggest that the primary target organ for DMT is the urinary tract with metabolism of DMT to TPA and formation of bladder calculi. This effect is reported in some but not all studies. Kidney weight reductions were observed in a 2-generation study. However, in the absence of histopathological changes in the kidney, this effect is considered to represent physiological adaptation (Jacobsen, 2005). Liver effects (diffuse hepatocellular swelling) were also reported in a 13-week dietary study but not in any other studies.

In a 14-day dietary study in rats, a NOAEL was determined as 660 mg/kg bw/d (males) and 1277 mg/kg bw/d (females) based on decreased body weights at higher doses (Chin et al., 1981). In this study, a NOAEL for induction of urinary calculi was 1320 mg/kg bw/d (males) and 1790 mg/kg bw/d (females). In contrast, in a 96-day dietary study in rats, a NOAEL of 313 mg/kg bw/d was determined based on decreased body weight gains at 636 mg/kg bw/d. In this study, there was no evidence of urinary calculi.

Therefore, an overall conservative LOAEL for reduced body weight gains was determined at 636 mg/kg bw/d from a 96-day rat study.

Inhalation repeat dose studies report no toxicological effects, including no urinary effects. Only mild and transient nose rubbing, preening, and blinking were observed in a 3-month inhalation study. A NOAEC of 86.4 mg/m³ was determined for inhalation exposure.

4.6 Genetic toxicity

Previous evaluations

The following genotoxicity studies were summarised from OECD (2001).

Test	Test system (species/strain)	Dose	Metabolic activation	Result	Reference
<i>In Vitro</i>					
Reverse mutation	<i>S. typhimurium</i> TA100, TA1535, TA98, TA1537	3.3-33 µg/plate	± S9	negative	Zeiger et al., 1982*
	<i>S. typhimurium</i> TA100, TA98	Unknown	± S9	negative	Kozumbo et al., 1982*
	<i>S. typhimurium</i> TA100, TA1535, TA98, TA1537, TA1538, TA102	0.5-5000 µg/plate	± S9	negative	Monarca et al., 1991*
	<i>Photobacterium phosphorium</i>	Unknown	± S9	negative	Elmore & Fitzgerald, 1990*

(table continued over page)

Test	Test system (species/strain)	Dose	Metabolic activation	Result	Reference
<i>In Vitro (cont.)</i>					
Unscheduled DNA synthesis	Rat hepatocytes	0, 727.5, 1455 or 2910 µg/mL	+ S9	negative	Monarca et al., 1991*
	SV40-transformed Chinese hamster cell line (CO60 cells)	0, 727.5, 1455 or 2910 µg/mL	- S9	negative	
	Human Hela cells	0, 5, 50 or 500 5000 µg/mL	± S9	negative	
	Hamster/Syrian (embryo cells)	0, 2.5, 5 or 10 µg/mL	- S9	negative	
Cytogenetic assay	Human lymphocytes	50-500 µg/mL	- S9	negative	Monarca et al., 1991*

Test	Test system (species/strain)	Dose	Metabolic activation	Result	Reference
Chromosomal aberration and sister chromatid exchange	Chinese hamster ovary cells (CHO)	10 µg/mL	± S9	negative	Loveday et al., 1990*
Mouse lymphoma mutation assay	L5178Y mouse lymphoma cells	0-100 µg/mL	± S9	negative	Myhr & Caspary, 1991*
<i>In Vivo</i>					
Sex-linked recessive lethality test	<i>Drosophila melanogaster</i>	1000 ppm (diet) or 400 ppm (injection) for 3 days	NA	negative	Foureman et al., 1994*
Micronucleus assay	B6C3F1 mice (bone marrow)	438-1750 mg/kg bw (intraperitoneal injection over 3 days)	NA	negative	Shelby*et al., 1993*
Micronucleus assay	(C57B1/6j x CBA)F1 mice (bone marrow)	0.20-1 mmol/kg bw (intraperitoneal injection)	NA	equivocal	Goncharova et al., 1988*

Data not reported in previous evaluations

No data.

Conclusion

The results from bacterial mutagenicity assays (Ames) all indicated that DMT is not mutagenic (Zeiger et al., 1982*; Kozumbo et al., 1982*; Monarca et al., 1991*; Elmore & Fitzgerald, 1990*). Negative results were also seen in in vitro DNA single-strand breakage assays, unscheduled DNA synthesis test, sister chromatid exchange assay, and in an assay evaluating the formation of micronuclei (Monarca et al., 1991*; Loveday et al., 1990*; Myhr & Caspary, 1991*). These studies utilised mouse hepatocytes, Chinese hamster embryo cells, HeLa cells, and human lymphocytes. Negative in vivo studies included a sex-linked recessive lethality test in *Drosophila* and one of two micronuclei studies in mice (Foureman et al., 1994*; Shelby et al., 1993*). The significance of an increase in micronuclei formation in one of these two in vivo tests is questionable due to many irregularities in the study's methodology as well as evidence of toxicity from the DMSO vehicle (Goncharova et al., 1988*). Furthermore, dose levels tested were much lower than those used in the other mouse micronuclei study (Shelby et al., 1993*) which showed a negative response.

On a weight-of-evidence basis, DMT is considered to be non-genotoxic.

4.7 Carcinogenicity

Previous evaluations

DMT was evaluated for carcinogenic potential in a 2-year dietary study (NCI, 1979*). F344 rats and B6C3F1 mice (50/sex/dose) were administered DMT via diet at doses of 0, 0.25 or 0.5% (0, 2500, 5000 ppm). These exposure levels did not affect body weight, survival or induce any clinical signs of toxicity. A statistically significant increase in the incidence of pulmonary tumours was observed in male mice. However, this was not considered to be treatment-related because the incidence observed fell within the historical control range for male B6C3F1 mice. No increases in any other tumour types were noted in either species.

In another study of a DMT metabolite, Wistar rats were given TPA in the diet at 0, 1, 2 and 5% (0, 10000, 20000 or 50000 ppm) for 2 years (Gross, 1974*, cited in Heck & Tyl, 1985). Bladder calculi were seen in both sexes only at the highest dose (42/47 and 39/42 for males and females, respectively). Bladder and ureteral neoplasms occurred in 21/37 males and 21/34 females at 5%, in 1/48 male and 2/47 females at 2% and in 1/43 males at 1%. No urinary tract neoplasms were detected in low dose females or in controls. Pituitary and thyroid tumours were noted amongst both treatment and control groups. These were not related to TPA dose. The investigator concluded that the presence of stones causes changes in the epithelium of the urinary tract ranging from hyperplasia, papilloma, squamous metaplasia to transitional cell tumours and squamous cell carcinoma. No other details were provided.

Data not reported in previous evaluations

No data.

Conclusion

A 2-year carcinogenicity study of DMT showed no treatment-related carcinogenic effects. Available data indicate that DMT is not carcinogenic in rats and mice. A 2-year carcinogenicity study of a DMT metabolite, TPA, reported that high doses (up to 50000 ppm) were associated with bladder stone formation, urinary tract neoplasia and carcinoma.

4.8 Reproductive toxicity

Traditional hazard assessments consider effects on fertility separate from developmental toxicity. Fertility is tested by exposing sexually mature adults to a chemical and examining the effects on reproductive capacity. Developmental toxicity is studied by exposing pregnant dams and looking for effects in the foetuses. Chemicals that affect the developing reproductive system following prenatal exposure may also affect sexual maturation or functional reproductive disorders that are only apparent at maturity. Developmental toxicity can therefore lead to effects on fertility and the two endpoints cannot be clearly distinguished.

In this hazard assessment, data are presented on the basis of test procedure. The effects on fertility and development are then discussed separately.

4.8.1 Repeat dose toxicity studies

Data not reported in previous evaluations

To investigate the effects of TPA on testicular function, male Sprague-Dawley rats (5 weeks old, 10 animals/dose) were given TPA in diet at 0, 0.2, 1 and 5% for 90 days (Cui et al., 2004). Mean TPA intakes calculated from food consumption were 0, 189, 1177 and 5818 mg/kg bw/d, respectively. Testicular functions were assessed by histology, testicular sperm head counts, daily sperm production, sperm motility, biochemical indices (marker testicular enzymes), and serum testosterone.

TPA caused no effects on food consumption or body weight of the rats. Serum testosterone concentrations and testes weight (relative or absolute) were unaffected. However, damage to spermatogenic cells and Sertoli cells at the 5% dose was reported. Decreases in testicular sperm head counts, daily sperm production and sorbitol dehydrogenase activity were also reported at this dose. Several characteristics of sperm motility such as VSL (straight line velocity), LIN (linearity), and STR (straightness) were reduced in treated groups in a dose-dependent manner. Mating performances were not determined.

4.8.2 Prenatal developmental toxicity studies

Previous evaluations

Exposure to 1000 mg/kg bw/d DMT via gavage from GD 7-16 in Wistar rats (sacrificed on day 21) had no effect on development (Hoechst, 1986*). Inhalation exposure in thirty rats to DMT at 1 mg/m³ throughout gestation had no effect on development (Krotov & Chebotar, 1972*). No further details were available for these studies.

Inhalation exposures to TPA at levels of 1, 5, and 10 mg/m³ (6 hours/day) from GD 6-15 in Sprague-Dawley rats had no effect on litter weights, pup viability or number of foetal malformations (Amoco Corporation, 1989*; Ryan et al., 1990*). A slight increase in rib anomalies was noted in the middle dose group, but rib anomalies were not considered to be an indicator of teratogenesis because they were common variations, not elevated in a dose-related manner, and within the range of historical controls.

4.8.3 One/two generation reproductive toxicity studies

Previous evaluations

Reproductive toxicity was evaluated in a one-generation reproduction study by exposing male Long Evans rats (30 rats/dose) to diets containing 0.25, 0.50 or 1.0% DMT for 115 days (Krasavage et al., 1973). These males were then mated with females that had been on the DMT diet for 6 days. After mating, females remained on the diet throughout pregnancy and lactation. No signs of toxicity were observed in either male or female parental animals. No effects on fertility parameters were noted. Pups born to parents fed 0.5 and 1.0% DMT had significantly lower average body weights at weaning compared to the controls. This effect on the offspring was likely due to exposure to DMT through lactation and having access to the mother's food. The NOAEL was 1.0% (636 mg/kg bw/d – the highest dose tested) for systemic and fertility effects in parents and 0.25% (152 mg/kg bw/d) for their offspring.

In a one-generation reproduction study, no adverse effects on fertility were noted in adult CD or Wistar rats fed with 0, 0.03, 0.125, 0.5, 2 or 5% TPA (a major metabolite of DMT) in the diet for 90 days prior to mating, through the mating period and continuing through gestation and lactation (CIIT, 1982*). Slight strain differences were noted. Reduced bodyweight gains were observed in CD rats at 2% and 5% and Wistar rats at 5%. There was no effect in either strain on fertility and litter size. There were increased postnatal deaths on day 1 and decreased pup survivability to day 21 at 2% and 5% in Wistar rats and at 5% for CD rats. Several litters of Wistar pups were lost to dams suffering obvious signs of maternal toxicity. Unscheduled deaths occurred during the post-weaning period in the 5% groups and was associated with a very high incidence of renal and bladder calculi in both rat strains. Weanling animals exhibited a higher incidence of calculi compared to adults consuming the same dietary level of TPA. The NOAEL for maternal and developmental toxicity was 0.5% (240-307 mg/kg bw/d), while the NOAEL for fertility effects was 5.0% (2480-3018 mg/kg bw/d).

Data not reported in previous evaluations

A 2-generation reproduction toxicity study conforming to OECD Test guideline 416 and GLP was conducted in rats to investigate the effect of diets containing TPA on fertility and reproductive performance. Twenty-six animals/group/sex were fed diets containing 0, 1000, 5000 or 20000 ppm TPA (approximately 98 – 261, 492 – 1218, 2011 - 5072 mg/kg bw/d). After 10 weeks, the animals (F0) were mated and allowed to rear the ensuing F1 litters to weaning. Breeding was repeated with the F₁ parents to produce F2 litters after a 10-week pre-mating period. Animals were fed continuously throughout the study (Milburn, 2003).

Reductions in bodyweight generally occurred when offspring started to consume solid diet (from day 15 post partum), and this was considered to be a direct effect of the test material on the pups rather than related to developmental toxicity. Because of the larger litter size at the 20000 ppm group in the F2 generation, pup bodyweights were also lower than the controls from parturition.

There were no effects on smear cycle and pattern, pre-coital interval, gestation length, proportion of successful matings, pups live born, pup survival, pup sex distribution or pup clinical observations in any dosed group. Sperm motility, sperm number and sperm morphology were also not affected by treatment with TPA.

There were no overt changes in the developmental landmarks in any pups examined in either generation. Non-treatment related observations related to reduced body weight include: delay in vaginal opening (1.6 days) in F1 females at 20000 ppm group; preputial separation in F1 males in the 5000 and 20000 ppm groups (0.8 and 1.6 days, respectively); and reduced AGD in females in the 20000ppm group. There was no evidence of a decrease in AGD in male pups. Differences in development landmarks were regarded as a secondary consequence of the reduction in body weights.

The no observed adverse effect level (NOAEL) for effects on fertility and development was 20000 ppm, the highest dose tested in the study.

Conclusion

Fertility effects

DMT in the diet up to 1% (636 mg/kg bw/d) had no effect on reproduction (Krasavage et al., 1973).

In a one-generation reproduction feeding study on TPA, no adverse effects on fertility were noted in adult rats fed up to 5% TPA (CIIT, 1982*). The NOAEL for fertility effects was >5.0% (2480-3018 mg/kg bw/d).

Similarly, in a 2-generation study in rats conducted according to OECD test guidelines and GLP, no effects on vaginal smears, reproductive performance, sperm motility, sperm number and sperm morphology were reported at 20000 ppm (approximately 2011-5072 mg/kg bw/d). The NOAEL for fertility in this study was 20000 ppm, the highest dose tested (Milburn, 2003).

In contrast, a subsequent study (Cui et al., 2004) of repeated doses of TPA on testicular function showed no effects on serum testosterone or testes weight but sperm head counts and daily sperm production decreased significantly at 5% (5818 mg/kg bw/d). In addition, several sperm motility parameters were also reduced significantly in all treated groups.

Developmental toxicity

DMT in the diet was associated with decreased pup weight at weaning but this effect was likely due to direct exposure to DMT through lactation and having access to the mother's food (Krasavage et al., 1973). The NOAEL for developmental effects was 152 mg/kg bw/d and the LOAEL 313 mg/kg bw/d, based on decreased body weight at weaning. The NOAEC in a developmental study for DMT was 1 mg/m³.

The NOAEL for oral exposure to TPA was at 0.5% (240-307 mg/kg bw/d) based on increased postnatal deaths and decreased pup survivability at higher doses (CIIT, 1982*). The adverse effects observed in the offspring in this study appear to be the result of maternal toxicity and the formation of renal and bladder calculi found in the weanling animals. The NOAEC in a developmental study for TPA was 10 mg/m³.

In a 2-generation study, there were no observed treatment-related changes in developmental landmarks (nipple/areolae development in males, preputial separation, vaginal opening and AGD) in any pups examined in either generation. Vaginal opening was slightly delayed (1.6 days) in F1 females at 20000 ppm group. Preputial separation was delayed in F1 males in the 5000 and 20000 ppm groups (0.8 and 1.6 days, respectively). However, the differences in development landmarks were regarded as a secondary consequence of the reduction in body weights. AGD was reduced in females in the 20000 ppm group, which is also considered to be a direct consequence of the slightly smaller size of animals at birth. There was no evidence of a decrease in AGD of male pups. The NOAEL for development effects was 20000 ppm (approximately 2011 - 5072 mg/kg bw/d), the highest dose tested in the study (Milburn, 2003).

5. Hazard Characterisation

Toxicity data for DMT were available for the majority of health endpoints. Additional data from terephthalic acid (TPA) are presented as TPA is a major metabolite of DMT.

DMT was readily absorbed after oral administration. Elimination was rapid with urine being the major route of excretion. There was no evidence of accumulation in tissues after multiple doses. The hydrolysis product, TPA, was the only metabolite detected in the urine in rats, while urinary metabolites in mice consisted of monomethyl terephthalate (70%) and TPA (30%).

DMT has low acute oral, dermal and inhalational toxicity. The acute oral LD₅₀ value in rats is 4390 – >6590 mg/kg bw, the dermal LD₅₀ in guinea pigs is >5000 mg/kg bw, and the inhalational LC₅₀ in rats is >6 mg/L.

DMT causes minimal skin and eye irritant effects in animals, and did not induce skin sensitisation in guinea pigs.

DMT does not appear to be mutagenic or genotoxic, nor was it deemed to be carcinogenic based on a 2-year feeding study.

For repeat dose toxicity, a conservative NOAEL based on reduced body weight gains in a rat 96-day dietary study was determined at 313 mg/kg bw/d. The target organ identified in oral exposure studies was the urinary tract with effects attributable to metabolism of DMT to TPA and the subsequent formation of bladder crystals or calculi primarily composed of Ca-TPA. In rats, bladder calculi formation occurred at a minimum DMT exposure length of 14 days at a dietary concentration of 1.5% (1890 mg/kg bw/d) for male rats and 2% (2290 mg/kg bw/d) for female rats.

Following 90-days of exposure via the inhalation route, a NOAEL of 86.4 mg/m³ was determined. Clinical effects at this dose, such as nose rubbing, preening, and blinking, were considered mild and transient. No urinary effects were observed following inhalation exposure.

In a one-generation reproduction study, DMT in the diet up to 1% (636 mg/kg bw/d) had no effect on reproduction although pup weights were reduced at weaning (Krasavage et al., 1973). There was no evidence of fertility effects, and hence the NOAEL for fertility was 636 mg/kg bw/d, the highest dose tested. In the same study, the NOAEL for developmental effects was 152 mg/kg bw/d, with a LOAEL of 313 mg/kg bw/d based on reduced pup weights at weaning, which was probably a lactational effect. The NOAEC in an inhalation developmental study for DMT was 1 mg/m³ and NOAEL in an oral developmental study was 1000 mg/kg.

The DMT metabolite, TPA, was also examined in reproduction studies. In a one-generation reproduction feeding study, no adverse effects on fertility were noted in adult rats fed up to 5% TPA (2480-3018 mg/kg bw/d). While the NOAEL for fertility effects was 5.0% (2480-3018 mg/kg bw/d), the NOAEL for maternal toxicity was 0.5%. The NOAEL for developmental toxicity was 0.5% (240-307 mg/kg bw/d), based on increased postnatal deaths and decreased pup survivability at higher doses. The adverse effects observed in the offspring in this study appear to be the result of

maternal toxicity and the formation of renal and bladder calculi in the weanling animals.

In a 2-generation study, rats treated with TPA showed no effects on smear cycle and pattern, pre-coital interval, gestation length, proportion of successful matings, pups live born, pup survival, pup sex distribution or pup clinical observations in any dosed group. Sperm motility, sperm number and sperm morphology were also not affected. There were no overt changes in the developmental landmarks in any pups examined in either generation. Delays in vaginal opening and preputial separation and reduced AGD in females in some animals were regarded as non-treatment related and linked to reductions in body weight. AGD were not decreased in male pups. Reductions in bodyweight generally occurred when the offspring started to consume solid diet (from day 15 post partum), and were considered to be a direct effect of the test material on the pups rather than related to developmental toxicity (Milburn, 2003). Because of the larger litter size at the 20000 ppm group in F2 generation, the pup bodyweights were also lower than controls from parturition. The NOAEL for effects on both fertility and development was 20000 ppm, (2011-5072 mg/kg bw/d) the highest dose tested in the study.

In an inhalation development study, the NOAEC for TPA was 10 mg/m³.

In contrast, a single study showed no effects on serum testosterone or testes weight but significant decreases in sperm head counts and daily sperm production in rats fed 5% TPA in the diet (5818 mg/kg bw/d). In addition, several sperm motility parameters also reduced significantly in all treated groups including at the lowest dose of 0.2%. Reproductive performance was not measured in this study.

With respect to fertility effects, a more recent OECD test guideline 2-generation reproduction study of di-2-ethylhexyl terephthalate (DEHT), also known to be metabolized to TPA in rats, also showed no effects on mating and fertility indices, oestrous cycle lengths, pre-coital intervals, gestation lengths, gender ratios, live litter sizes or postnatal survival for either F0 or F1 generations up to the highest dose tested of 1% in the diet (F1: 893-1549 mg/kg bw/d) (Faber et al., 2007). Therefore, overall, weight-of-evidence from available studies indicates that DMT or its metabolite TPA do not affect fertility, at doses up to approximately 5000 mg/kg bw/d. In addition, no developmental effects unrelated to maternal toxicity were observed with DMT in oral and inhalation studies.

6. Human Health Hazard Summary Table

Phthalate	Acute Toxicity	Irritation & Sensitisation	Repeated Dose Toxicity	Genetic Toxicity	Carcinogenicity	Fertility	Developmental Toxicity
Dimethyl terephthalate (DMT)	<p>Oral Rat: LD50 = 4390 – > 6590 mg/kg bw</p> <p>Dermal Guinea pig: LD50 > 5000 mg/kg bw</p> <p>Inhalation Rat: LC50 >6 mg/L</p>	<p>Skin irritation: Minimal effects</p> <p>Eye irritation: Minimal effects</p> <p>Respiratory irritation: No data</p> <p>Skin sensitisation: Negative</p>	<p>Rat: Oral NOAEL = 313 mg/kg bw/d LOAEL = 636 mg/kg bw/d, ↓ body weight gains.</p> <p>Inhalation NOAEL = 1 mg/m³</p> <p>High doses: Bladder calculi and urothelial hyperplasia.</p>	<p><i>In vitro</i> Negative in Ames test, DNA damage, sister chromatid exchange and micronuclei assays</p> <p><i>In vivo</i> Negative in sex-linked recessive lethality and micronuclei tests</p>	<p>Rat, mouse: no treatment-related carcinogenicity.</p> <p>NOAEL = 300 mg/kg bw/d</p>	<p>Two-generation study Rat: NOAEL = 5072 mg/kg bw/d (TPA) LOAEL = not established</p>	<p>Two-generation study Rat: NOAEL = 5072 mg/kg bw/d (TPA) LOAEL = not established</p>

Page intentionally blank for double-sided printing.

References

- Amoco Corporation (1989) A segment II inhalation teratology study of terephthalic acid in rats. IIT Research Institute, IITRI Study no. 1448.
- Anonymous (1986) Prehled Prumyslove Toxikol Org Latky, p 386 RTECS (1988). Cited in Du Pont report.
- Chin TY, Tyl RW, Popp JA, & Heck HD (1981) Chemical urolithiasis. 1. Characteristics of bladder stone induction by terephthalic acid and dimethyl terephthalate in weanling Fischer-344 rats. *Toxicol Appl Pharmacol*, 58:307-21.
- CIIT (1982) A ninety-day study of terephthalic acid-induced urolithiasis and reproductive performance in Wistar and CD rats. [Report prepared for the Chemical Industry Institute of Toxicology by IIT Research Institute, CIIT Docket No. 11622]. Research Triangle Park, NC, Chemical Industry Institute of Toxicology.
- Cui L, Dai G, Xu L, Wang S, Song L, Zhao R, Xiao H, Zhou J, & Wang X (2004) Effect of oral administration of terephthalic acid on testicular functions of rats. *Toxicology*, 201:59-66.
- Davidenko AV et al. (1984) *Biol.Nauki (Moskov)* 1:31-34 (CA 100:11602v).
- Eastman Kodak Co. (1957). Unpublished data.
- Eastman Kodak Co. (1963). Unpublished data.
- Elmore E & Fitzgerald MP (1990) Evaluation of the bioluminescence assays as screens for genotoxic chemicals. *Prog Clin Biol Res*, 340D:379-87.
- Faber WD, Deyo JA, Stump, DG and Ruble K (2007) Two-generation reproduction study of di-2-ethylhexyl terephthalate in Crl:CD rats. *Birth Defects Research* 80: 69-81
- Foureman P, Mason JM, Valencia R, & Zimmering S (1994) Chemical mutagenesis testing in *Drosophila*. X. Results of 70 coded chemicals tested for the National Toxicology Program. *Environ Mol Mutagen*, 23:208-27.
- Goncharova RI, Zabrejko S, Kozachenko VI, & Pashin YuV (1988) Mutagenic effects of dimethyl terephthalate on mouse somatic cells in vivo. *Mutat Res*, 204:703-9.
- Gross J (1974) The effects of prolonged feeding of terephthalic acid (TPA) to rats. Project FG-Is-175. Agricultural Research Service, US Department of Agriculture, Washington, DC. Cited in Heck & Tyl (1985).
- Heck HD & Tyl RW (1985) The induction of bladder stones by terephthalic acid, dimethyl terephthalate, and melamine (2,4,6-triamino-s-triazine) and its relevance to risk assessment. *Regul Toxicol Pharmacol*, 5:294-313.
- Heck Hd'A (1980) Metabolism of dimethyl terephthalate (DMT) in F-344 rats and B6C3F1 mice. Abstracts, 19th Annual Meeting of the Society of Toxicology, A81 (Abstract 242), Washington DC, March 9-13.
- Hoechst AG (1986) Dimethyl terephthalate, investigation of embryotoxic action in Wistar rats on oral administration. Unpublished report no. 86.0859. Commissioned by the employment accident insurance fund of the chemical industry. BG Chemie Toxicological Evaluations 1 Potential Health Hazards of Existing Chemicals, Springer-Verlag.
- Jacobsen M (2005) Terephthalic acid: Multigeneration reproduction toxicity study in rats. Pathology comment on the renal findings from study RR0915. Unpublished report. Central Toxicology Laboratory, Macclesfield, UK.

- Kamal'Dinova ZM et al. (1962) Prom Tokiol I Klinkia Prof Zabol Khim Etiol Sb, 159-160 (CA 61: 11230g). Cited in OECD SIAR Dimethyl Terephthalate, 2001.
- Kozumbo WJ, Kroll R, & Rubin RJ (1982) Assessment of the mutagenicity of phthalate esters. *Environ Health Perspect*, 45:103-9.
- Krasavage WJ, Yanno FJ, & Terhaar CJ (1973) Dimethyl terephthalate (DMT): acute toxicity, subacute feeding and inhalation studies in male rats. *Am Ind Hyg Assoc J*, 34:455-62.
- Krotov IuA & Chebotar NA (1972) Study of the embryotoxic and teratogenic effect of several industrial substances formed during the production of dimethylterephthalate [in Russian]. *Gig Tr Prof Zabol*, 16:40-3 (CA 77:97441V) (Translation in J-927).
- Lewis TR et al. (1982) *The Toxicologist* 2:1 (Abstract 25).
- Loveday KS, Anderson BE, Resnick MA, & Zeiger E (1990) Chromosome aberration and sister chromatid exchange tests in Chinese hamster ovary cells in vitro. V: Results with 46 chemicals. *Environ Mol Mutagen*, 16:272-303.
- Marhold JV (1972) Spornik Vysledku Tokikologickho Vystreni Latek A Priparavki, p47 in RTECS (1976)
- Milburn GM (2003) Terephthalic acid: Multigeneration reproduction toxicity study in rats. Unpublished report no. CTL/RR0915. Central Toxicology Laboratory, Macclesfield, UK.
- Moffitt AE Jr, Clary JJ, Lewis TR, Blanck MD, & Perone VB (1975) Absorption, distribution and excretion of terephthalic acid and dimethyl terephthalate. *Am Ind Hyg Assoc J*, 36:633-41.
- Monarca S, Pool-Zobel BL, Rizzi R, Klein P, Schmezer P, Piatti E, Pasquini R, De Fusco R, & Biscardi D (1991) In vitro genotoxicity of dimethyl terephthalate. *Mutat Res*, 262:85-92.
- Myhr BC & Caspary WJ (1991) Chemical mutagenesis at the thymidine kinase locus in L5178Y mouse lymphoma cells: results for 31 coded compounds in the National Toxicology Program. *Environ Mol Mutagen*, 18:51-83.
- NCI (1979) Bioassay of dimethyl terephthalate for possible carcinogenicity, CAS no. 120-61-6. Bethesda, Maryland, National Cancer Institute Carcinogenesis Technical Report Series no. 121 (NCI-CG-TR-121). NIH Publication no. 79-1376.
- OECD (2001) SIDS Initial Assessment Report for SIAM 11: Dimethyl terephthalate. Organisation for Economic Cooperation and Development, USA, 23-26 January 2001.
- Patty's Industrial Hygiene and Toxicology (1981) 3rd revised Edition, Volume IIA: 2344-2345, 2348-2349, 2352.
- Ryan BM, Hatoum NS, & Jernigan JD (1990) A segment II inhalation teratology study of terephthalic acid in rats. *Toxicologist*, 10:40.
- Sanina YP & Kocketkova TA (1963) Toksikol. Novykh. Prom. Khim. Veshchestv, (5):107-123 (CA 61:6250f).
- Shelby MD, Erexson GL, Hook GJ, & Tice RR (1993) Evaluation of a three-exposure mouse bone marrow micronucleus protocol: results with 49 chemicals. *Environ Mol Mutagen*, 21:160-79.
- Vogin EE (1972) Subacute feeding studies (13-week) in rats with dimethylterephthalate (DMT), isophthalic acid (IA), and terephthalic acid (TA) (unpublished data). Food and Drug Research Laboratories, Inc., Maspeth, NY. Cited in Heck & Tyl (1985).
- Zeiger E, Haworth S, Speck W, Mortelmans K (1982) Phthalate ester testing in the National Toxicology Program's environmental mutagenesis test development program. *Environ Health Perspect*, 45:99-101.